

Statistical Analysis in Lexis Diagrams: Age-Period-Cohort models — and some cousins

Bendix Carstensen Steno Diabetes Center Copenhagen, Gentofte, Denmark
<http://BendixCarstensen.com>

European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

From /home/bendix/teach/APC/courses/EDSD_2020/slides/s1.tex

Wednesday 27th May, 2020, 07:43

1 / 333

About the practicals

- ▶ You should use your preferred **R**-environment.
- ▶ Epi-package for **R** is needed, check that you have version 2.40
- ▶ Data are all on the course website.
- ▶ Try to make a text version of the answers to the exercises — it is more rewarding than just looking at output. The latter is soon forgotten — Rmd is a possibility.
- ▶ An opportunity to learn emacs, ESS and Sweave?

Introduction (intro)

5 / 333

Introduction

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

intro

Rates and Survival

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

surv-rate

Welcome

- ▶ Purpose of the course:
 - ▶ knowledge about APC-models
 - ▶ technical knowledge of handling them
 - ▶ insight in the basic concepts of analysis of rates
 - ▶ handling observation in the Lexis diagram
- ▶ Remedies of the course:
 - ▶ Lectures with handouts (BxC)
 - ▶ Practicals with suggested solutions (BxC)
 - ▶ Assignment for Thursday

Introduction (intro)

2 / 333

Survival data

- ▶ Persons enter the study at some date.
- ▶ Persons exit at a later date, either dead or alive.
- ▶ Observation:
 - ▶ Actual time span to death ("event")
 - ▶ ... or ...
 - ▶ Some time alive ("at least this long")

Rates and Survival (surv-rate)

6 / 333

Scope of the course

- ▶ Rates as observed in populations — disease registers for example.
- ▶ Understanding of survival analysis (statistical analysis of rates) — this is the content of much of the first day.
- ▶ Besides concepts, practical understanding of the actual computations (in **R**) are emphasized.
- ▶ There is a section in the practicals: "Basic concepts of rates and survival" — read it; use it as reference.
- ▶ If you are not quite familiar with matrix algebra in **R**, there is an intro on the course homepage.

Introduction (intro)

3 / 333

Examples of time-to-event measurements

- ▶ Time from diagnosis of cancer to death.
- ▶ Time from randomization to death in a cancer clinical trial
- ▶ Time from HIV infection to AIDS.
- ▶ Time from marriage to 1st child birth.
- ▶ Time from marriage to divorce.
- ▶ Time from jail release to re-offending

Rates and Survival (surv-rate)

7 / 333

About the lectures

- ▶ Please interrupt: Most likely I did a mistake or left out a crucial argument.
- ▶ The handouts are not perfect — please comment on them, prospective students would benefit from it.
- ▶ Time-schedule: Two lectures (≈ 2 hrs) one practical for you to do (≈ 1 hr)
- ▶ Practical will be recapped next morning


Introduction (intro)

4 / 333

Each line a person

Each blob a death

Study ended at 31 Dec. 2003




Rates and Survival (surv-rate)

8 / 333

Ordered by date of entry

Most likely the order in your database.




Survival after Cervix cancer

Year	Stage I		Stage II	
	N	L	N	L
1	110	5	234	24
2	100	7	207	27
3	86	7	169	31
4	72	3	129	17
5	61	0	105	7
6	54	2	85	6
7	42	3	73	5
8	33	0	62	3
9	28	0	49	2
10	24	1	34	4

Estimated risk in year 1 for Stage I women is $5/107.5 = 0.0465$

Estimated 1 year survival is $1 - 0.0465 = 0.9535$ — Life-table estimator.

Timescale changed to "Time since diagnosis".



Survival function

Persons enter at time 0:

- Date of birth
- Date of randomization
- Date of diagnosis.


How **long** they survive, survival time T — a stochastic variable.

Distribution is characterized by the survival function:

$$S(t) = P\{\text{survival at least till } t\}$$

$$= P\{T > t\} = 1 - P\{T \leq t\} = 1 - F(t)$$

Patients ordered by survival time.



Intensity or rate

$$\lambda(t) = P\{\text{event in } (t, t+h] \mid \text{alive at } t\} / h$$

$$= \frac{F(t+h) - F(t)}{S(t) \times h}$$


$$= - \frac{S(t+h) - S(t)}{S(t)h} \xrightarrow{h \rightarrow 0} - \frac{d \log S(t)}{dt}$$

This is the **intensity** or **hazard function** for the distribution.

Characterizes the survival distribution as does f or F .

Theoretical counterpart of a **rate**.


Survival times grouped into bands of survival.




Empirical rates for individuals

- At the **individual** level we introduce the **empirical rate**: (d, y) ,
 - no. of events ($d \in \{0, 1\}$) during y risk time
- Each person may contribute several empirical (d, y)
- Empirical rates are **responses** in survival analysis
- The timescale is a **covariate**:
 - that varies between empirical rates from one individual: Age, calendar time, time since diagnosis
- Do not confuse timescale with y — risk time (called exposure in demography)
 - a **difference** between two points on **any** timescale


Patients ordered by survival status within each band.



Empirical rates by calendar time.



Empirical rates by time since diagnosis.



Likelihood for rates

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

Two timescales

Note that we actually have two timescales:

- ▶ Time since diagnosis (*i.e.* since entry into the study)
- ▶ Calendar time.

These can be shown **simultaneously** in a Lexis diagram.

Likelihood contribution from one person

The likelihood from several empirical rates from one individual is a product of conditional probabilities:

$$P\{\text{event at } t_4 | \text{alive at } t_0\} = P\{\text{event at } t_4 | \text{alive at } t_3\} \times P\{\text{survive } (t_2, t_3) | \text{alive at } t_2\} \times P\{\text{survive } (t_1, t_2) | \text{alive at } t_1\} \times P\{\text{survive } (t_0, t_1) | \text{alive at } t_0\}$$


Likelihood contribution from one individual is a **product** of terms.

Each term refers to one empirical rate (d, y) with $y = t_{i+1} - t_i$ (mostly $d = 0$).

t_i is a **covariate**

Follow-up by calendar time *and* time since diagnosis:

A Lexis diagram!





Likelihood for an empirical rate

- ▶ Likelihood depends on **data** and the **model**
- ▶ Model: the rate (λ) is constant in the interval.
- ▶ The interval should be sufficiently small for this assumption to be reasonable.

$$L(\lambda | y, d) = P\{\text{survive } y\} \times P\{\text{event}\}^d = e^{-\lambda y} \times (\lambda dt)^d = \lambda^d e^{-\lambda y}$$

$$\ell(\lambda | y, d) = d \log(\lambda) - \lambda y$$

Empirical rates by calendar time *and* time since diagnosis



Probability

$$P(d \text{ at } t_x | \text{entry } t_0)$$

$$= P(\text{surv } t_0 \rightarrow t_1 | \text{entry } t_0) \times P(\text{surv } t_1 \rightarrow t_2 | \text{entry } t_1) \times P(d \text{ at } t_x | \text{entry } t_2)$$


log-Likelihood

$$d \log(\lambda) - \lambda y$$

$$= 0 \log(\lambda) - \lambda y_1 + 0 \log(\lambda) - \lambda y_2 + d \log(\lambda) - \lambda y_3$$

So what's the purpose

- ▶ form the basis for statistical inference about occurrence rates:
- ▶ response: observed rates of events and person-time (d, y)
- ▶ covariates:
 - ▶ **A**: Age at follow-up
 - ▶ **P**: Period (date) of follow-up
 - ▶ **C**: (=P-A) Cohort (date of birth)



Probability


$$P(\text{surv } t_0 \rightarrow t_x | \text{entry } t_0)$$

$$= P(\text{surv } t_0 \rightarrow t_1 | \text{entry } t_0) \times P(\text{surv } t_1 \rightarrow t_2 | \text{entry } t_1) \times P(\text{surv } t_2 \rightarrow t_x | \text{entry } t_2)$$

log-Likelihood

$$0 \log(\lambda) - \lambda y$$

$$= 0 \log(\lambda) - \lambda y_1 + 0 \log(\lambda) - \lambda y_2 + 0 \log(\lambda) - \lambda y_3$$




Probability

$$\begin{aligned}
 &P(\text{event at } t_x | \text{entry } t_0) \\
 &= P(\text{surv } t_0 \rightarrow t_1 | \text{entry } t_0) \\
 &\times P(\text{surv } t_1 \rightarrow t_2 | \text{entry } t_1) \\
 &\times P(\text{event at } t_x | \text{entry } t_2)
 \end{aligned}$$

log-Likelihood

$$\begin{aligned}
 &1 \log(\lambda) - \lambda y \\
 &= 0 \log(\lambda) - \lambda y_1 \\
 &+ 0 \log(\lambda) - \lambda y_2 \\
 &+ 1 \log(\lambda) - \lambda y_3
 \end{aligned}$$

Log-likelihood ratio




Aim of dividing time into bands:

- ▶ Compute rates in different bands of:
 - ▶ age
 - ▶ calendar time
 - ▶ disease duration
 - ▶ ...
- ▶ Allow rates to vary along the timescale:

$$\begin{aligned}
 &0 \log(\lambda) - \lambda y_1 \\
 &+ 0 \log(\lambda) - \lambda y_2 \\
 &+ d \log(\lambda) - \lambda y_3
 \end{aligned}
 \rightarrow
 \begin{aligned}
 &0 \log(\lambda_1) - \lambda_1 y_1 \\
 &+ 0 \log(\lambda_2) - \lambda_2 y_2 \\
 &+ d \log(\lambda_3) - \lambda_3 y_3
 \end{aligned}$$

Log-likelihood ratio




Log-likelihood from more persons

- ▶ One person p , different times t : $\sum_t (d_{pt} \log(\lambda_t) - \lambda_t y_{pt})$
- ▶ More persons: $\sum_p \sum_t (d_{pt} \log(\lambda_t) - \lambda_t y_{pt})$
- ▶ Collect terms with identical values of λ_t :

$$\begin{aligned}
 \sum_t \sum_p (d_{pt} \log(\lambda_t) - \lambda_t y_{pt}) &= \sum_t ((\sum_p d_{pt}) \log(\lambda_t) - \lambda_t \sum_p y_{pt}) \\
 &= \sum_t (D_t \log(\lambda_t) - \lambda_t Y_t)
 \end{aligned}$$

- ▶ All events in interval t ("at" time t), D_t
- ▶ All exposure time in interval t ("at" time t), Y_t

Log-likelihood ratio




Likelihood example

- ▶ Assuming the rate (intensity) is constant, λ ,
- ▶ the probability of observing 7 deaths in the course of 500 person-years:


$$\begin{aligned}
 P\{D = 7, Y = 500 | \lambda\} &= \lambda^D e^{-\lambda Y} \times K \\
 &= \lambda^7 e^{-\lambda 500} \times K \\
 &= L(\lambda | \text{data})
 \end{aligned}$$

- ▶ Best guess of λ is where this function is as large as possible.
- ▶ Confidence interval is where it is not too far from the maximum


Log-likelihood ratio



Likelihood-ratio function



Log-likelihood ratio



Poisson likelihood

Log-likelihood from **follow-up** of **one individual**, p , in interval t :

$$\ell_{\text{FU}}(\lambda|d, y) = d_{pt} \log(\lambda(t)) - \lambda(t) y_{pt}, \quad t = 1, \dots, t_p$$

Log-likelihood from a **Poisson observation** d_{pt} with mean $\mu = \lambda(t) y_{pt}$:

$$\begin{aligned} \ell_{\text{Poisson}}(\lambda y|d) &= d_{pt} \log(\lambda(t) y_{pt}) - \lambda(t) y_{pt} \\ &= \ell_{\text{FU}}(\lambda|d, y) + d_{pt} \log(y_{pt}) \end{aligned}$$

Extra term does not depend on the rate parameter λ .

Confidence interval for a rate

A 95% confidence interval for the log of a rate is:

$$\hat{\theta} \pm 1.96/\sqrt{D} = \log(\lambda) \pm 1.96/\sqrt{D}$$

Take the exponential to get the confidence interval for the rate:

$$\lambda \times \underbrace{\exp(1.96/\sqrt{D})}_{\text{error factor, erf}}$$

Alternatively do the c.i. directly on the rate scale:

$$\lambda \pm 1.96\sqrt{D}/Y$$

Poisson likelihood

Log-likelihood contribution from **one** individual, p , say, is:

$$\ell_{\text{FU}}(\lambda|d, y) = \sum_t (d_{pt} \log(\lambda(t)) - \lambda(t) y_{pt})$$

- ▶ The terms in the sum are **not** independent,
 - ▶ but the log-likelihood is a **sum** of Poisson-like terms,
 - ▶ the **same** as a likelihood for **independent** Poisson variates, d_{pt}
 - ▶ with mean $\mu = \lambda_t y_{pt} \Leftrightarrow \log \mu = \log(\lambda_t) + \log(y_{pt})$
- ⇒ Analyze rates λ based on empirical rates (d, y) as a Poisson model for independent variates where:
- ▶ d_{pt} is the response variable.
 - ▶ $\log(y_{pt})$ is the offset variable.

Exercise

Suppose we have 17 deaths during 843.6 years of follow-up.

Calculate the mortality rate with a 95% c.i.

Likelihood for follow-up of many subjects

Adding empirical rates over the follow-up of persons:

$$D = \sum d \quad Y = \sum y \quad \Rightarrow \quad D \log(\lambda) - \lambda Y$$

- ▶ Persons are assumed independent
- ▶ Contribution from the same person are **conditionally** independent, hence give separate contributions to the log-likelihood.
- ▶ Follow-up **model** and Poisson **model** are different
- ▶ ... but the **likelihoods** are the same.

Rates with glm

```
> library(Epi)
> D <- 17
> Y <- 843.6/1000
> round( ci.exp( glm( D ~ 1, offset=log(Y), family=poisson ), 2 )
          exp(Est.)  2.5% 97.5%
(Intercept)  20.15 12.53 32.42
> round( ci.exp( glm( cbind(D,Y) ~ 1, family=poisreg ), 2 )
          exp(Est.)  2.5% 97.5%
(Intercept)  20.15 12.53 32.42
> round( ci.exp( glm( cbind(D,Y) ~ 1, family=poisreg(link="identity"),
+                 Exp=FALSE ), 2 )
          Estimate  2.5% 97.5%
(Intercept)  20.15 10.57 29.73
```

The log-likelihood is maximal for:

$$\frac{d\ell(\lambda)}{d\lambda} = \frac{D}{\lambda} - Y = 0 \quad \Leftrightarrow \quad \hat{\lambda} = \frac{D}{Y}$$

Information about the log-rate $\theta = \log(\lambda)$:

$$\ell(\theta|D, Y) = D\theta - e^\theta Y, \quad \ell'_\theta = D - e^\theta Y, \quad \ell''_\theta = -e^\theta Y$$

so $I(\hat{\theta}) = e^{\hat{\theta}} Y = \hat{\lambda} Y = D$, hence $\text{var}(\hat{\theta}) = 1/D$

Standard error of log-rate: $1/\sqrt{D}$.

Note that this only depends on the no. events, **not** on the follow-up time.

Ratio of two rates

If we have observations two rates λ_1 and λ_0 , based on (D_1, Y_1) and (D_0, Y_0) the variance of the log of the ratio of the rates, $\log(\text{RR})$, is:

$$\begin{aligned} \text{var}(\log(\text{RR})) &= \text{var}(\log(\lambda_1/\lambda_0)) \\ &= \text{var}(\log(\lambda_1)) + \text{var}(\log(\lambda_0)) \\ &= 1/D_1 + 1/D_0 \end{aligned}$$

As before, a 95% c.i. for the RR is then:

$$\text{RR} \times \underbrace{\exp\left(1.96\sqrt{\frac{1}{D_1} + \frac{1}{D_0}}\right)}_{\text{error factor}}$$

The log-likelihood is maximal for:

$$\frac{d\ell(\lambda)}{d\lambda} = \frac{D}{\lambda} - Y = 0 \quad \Leftrightarrow \quad \hat{\lambda} = \frac{D}{Y}$$

Information about the rate itself, λ :

$$\ell(\lambda|D, Y) = D \log(\lambda) - \lambda Y \quad \ell'_\lambda = \frac{D}{\lambda} - Y \quad \ell''_\lambda = -\frac{D}{\lambda^2}$$

so $I(\hat{\lambda}) = D/\hat{\lambda}^2 = Y^2/D$, hence $\text{var}(\hat{\lambda}) = D/Y^2$

Standard error of a rate: $\sqrt{D}/Y = \hat{\lambda}/\sqrt{D}$.

Exercise

Suppose we in group 0 have 17 deaths during 843.6 years of follow-up in one group, and in group 1 have 28 deaths during 632.3 years.

Calculate the rate-ratio between group 1 and 0 with a 95% c.i.

Rate-ratio with glm

```
> library(Epi)
> D <- c(17,28)
> Y <- c(843.6,632.3)/1000
> F <- factor(0:1)
> round( ci.exp( glm( cbind(D,Y) ~ F , family=poisreg ) , 2 )
      exp(Est.)  2.5% 97.5%
(Intercept)  20.15 12.53 32.42
F1           2.20  1.20  4.01

> round( ci.exp( glm( cbind(D,Y) ~ F - 1, family=poisreg ) , 2 )
      exp(Est.)  2.5% 97.5%
FO           20.15 12.53 32.42
F1           44.28 30.58 64.14
```

Likelihood for rates (likelihood)

52/ 333

Population life table, DK 1997–98

a	Men			Women		
	S(a)	λ(a)	E[l _{res} (a)]	S(a)	λ(a)	E[l _{res} (a)]
0	1.00000	567	73.68	1.00000	474	78.65
1	0.99433	67	73.10	0.99526	47	78.02
2	0.99366	38	72.15	0.99479	21	77.06
3	0.99329	25	71.18	0.99458	14	76.08
4	0.99304	25	70.19	0.99444	14	75.09
5	0.99279	21	69.21	0.99430	11	74.10
6	0.99258	17	68.23	0.99419	6	73.11
7	0.99242	14	67.24	0.99413	3	72.11
8	0.99227	15	66.25	0.99410	6	71.11
9	0.99213	14	65.26	0.99404	9	70.12
10	0.99199	17	64.26	0.99395	17	69.12
11	0.99181	19	63.28	0.99378	15	68.14
12	0.99162	16	62.29	0.99363	11	67.15
13	0.99147	18	61.30	0.99352	14	66.15
14	0.99129	25	60.31	0.99338	11	65.16
15	0.99104	45	59.32	0.99327	10	64.17
16	0.99059	50	58.35	0.99317	18	63.18
17	0.99009	52	57.38	0.99299	29	62.19
18	0.98957	85	56.41	0.99270	35	61.21
19	0.98873	79	55.46	0.99235	30	60.23
20	0.98795	70	54.50	0.99205	35	59.24
21	0.98726	71	53.54	0.99170	31	58.27

Lifetables (lifetable)

56/ 333

Rate-ratio and -difference with glm


```
> round( ci.exp( glm( cbind(D,Y) ~ F , family=poisreg ) , 2 )
      exp(Est.)  2.5% 97.5%
(Intercept)  20.15 12.53 32.42
F1           2.20  1.20  4.01

> round( ci.exp( glm( cbind(D,Y) ~ F , family=poisreg(link="identity"),
+               Exp=FALSE, 2 )
      Estimate  2.5% 97.5%
(Intercept)  20.15 10.57 29.73
F1           24.13  5.14 43.13

> round( ci.exp( glm( cbind(D,Y) ~ F - 1, family=poisreg(link="identity"),
+               Exp=FALSE, 2 )
      Estimate  2.5% 97.5%
FO           20.15 10.57 29.73
F1           44.28 27.88 60.69
```

Likelihood for rates (likelihood)

53/ 333



Lifetables (lifetable)

57/ 333


Lifetables

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

lifetable



Lifetables (lifetable)

58/ 333

The life table method

The simplest analysis is by the "life-table method":

interval	alive	dead	cens.	
i	n_i	d_i	l_i	p_i
1	77	5	2	$5/(77 - 2/2) = 0.066$
2	70	7	4	$7/(70 - 4/2) = 0.103$
3	59	8	1	$8/(59 - 1/2) = 0.137$

$$p_i = P\{\text{death in interval } i\} = 1 - d_i/(n_i - l_i/2)$$

$$S(t) = (1 - p_1) \times \dots \times (1 - p_t)$$

Lifetables (lifetable)

54/ 333

Practical

Based on the previous slides answer the following for both Danish and Swedish life tables:

- ▶ What is the doubling time for mortality?
- ▶ What is the rate-ratio between males and females?
- ▶ How much older should a woman be in order to have the same mortality as a man?

Lifetables (lifetable)

59/ 333

The life table method

The life-table method computes survival probabilities for each time interval, in demography normally one year.

The rate is the number of deaths d_i divided by the risk time $(n_i - d_i/2 - l_i/2) \times \ell_i$:

$$\lambda_i = \frac{d_i}{(n_i - d_i/2 - l_i/2) \times \ell_i}$$

and hence the death probability:

$$p_i = 1 - \exp(-\lambda_i \ell_i) = 1 - \exp\left(-\frac{d_i}{(n_i - d_i/2 - l_i/2)}\right)$$

The modified life-table estimator.

Lifetables (lifetable)

55/ 333


Denmark	Males	Females
$\log_2(\lambda(a))$	$-14.244 + 0.135 \text{ age}$	$-14.877 + 0.135 \text{ age}$
Doubling time	$1/0.135 = 7.41 \text{ years}$	
M/F rate-ratio	$2^{-14.244+14.877} = 2^{0.633} = 1.55$	
Age-difference	$(-14.244 + 14.877)/0.135 = 4.69 \text{ years}$	

Sweden:	Males	Females
$\log_2(\lambda(a))$	$-15.453 + 0.146 \text{ age}$	$-16.204 + 0.146 \text{ age}$
Doubling time	$1/0.146 = 6.85 \text{ years}$	
M/F rate-ratio	$2^{-15.453+16.204} = 2^{0.751} = 1.68$	
Age-difference	$(-15.453 + 16.204)/0.146 = 5.14 \text{ years}$	

Lifetables (lifetable)

60/ 333

Observations for the lifetable



Life table is based on person-years and deaths accumulated in a short period.

Age-specific rates — cross-sectional!

Survival function:

$$S(t) = e^{-\int_0^t \lambda(a) da} = e^{-\sum_0^t \lambda(a)}$$

— assumes stability of rates to be interpretable for actual persons.

Who needs the Cox-model anyway?

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:

Age-Period-Cohort models

— and some cousins

European Doctoral School of Demography,

Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

Life table approach

The observation of interest is **not** the survival time of the **individual**.

It is the **population** experience:

D : Deaths (events).

Y : Person-years (risk time).

The classical lifetable analysis compiles these for prespecified intervals of age, and computes age-specific mortality **rates**.

Data are collected cross-sectionally, but interpreted longitudinally.

A look at the Cox model


$$\lambda(t, x) = \lambda_0(t) \times \exp(x'\beta)$$

A model for the rate as a function of t and x .

Covariates:

- ▶ x
- ▶ t
- ▶ ... often the effect of t is ignored (forgotten?)
- ▶ *i.e.* left unreported

Rates vary over time:




The Cox-likelihood as profile likelihood

- ▶ One parameter per death time to describe the effect of time (i.e. the chosen timescale).


$$\log(\lambda(t, x_i)) = \log(\lambda_0(t)) + \underbrace{\beta_1 x_{1i} + \dots + \beta_p x_{pi}}_{\eta_i} = \alpha_t + \eta_i$$

- ▶ Profile likelihood:
 - ▶ Derive estimates of α_t as function of data and β s — assuming constant rate between death/censoring times
 - ▶ Insert in likelihood, now only a function of data and β s
 - ▶ This turns out to be Cox's partial likelihood
- ▶ Cumulative intensity ($\Lambda_0(t)$) obtained via the Breslow-estimator


Rates vary over time:



Mayo Clinic lung cancer data: 60 year old woman



Rates vary over time:



Splitting the dataset a priori

- ▶ The Poisson approach needs a dataset of empirical rates (d, y) with suitably small values of y .
- ▶ — each individual contributes many empirical rates
- ▶ (one per risk-set contribution in Cox-modeling)
- ▶ From each empirical rate we get:
 - ▶ Poisson-response d
 - ▶ Risk time $y \rightarrow \log(y)$ as offset
 - ▶ time scale covariates: current age, current date, ...
 - ▶ other covariates
- ▶ Contributions not independent, but likelihood is a product
- ▶ Same likelihood as for independent Poisson variates
- ▶ Poisson `glm` with spline/factor effect of time

Example: Mayo Clinic lung cancer

- ▶ Survival after lung cancer
- ▶ Covariates:
 - ▶ Age at diagnosis
 - ▶ Sex
 - ▶ Time since diagnosis
- ▶ Cox model
- ▶ Split data:
 - ▶ Poisson model, time as factor
 - ▶ Poisson model, time as spline

Who needs the Cox-model anyway? (RXCox)

68/ 333

Example: Mayo Clinic lung cancer IV

```
> length( coef(mLs.pois.fc) )
[1] 188

> t.kn <- c(0,25,100,500,1000)
> dim( Ns(Lung.s$tfe,knots=t.kn) )
[1] 20022 4


> system.time(
+ mLs.pois.sp <- glm( cbind(lex.Xst=="Dead",lex.dur) ~
+ Ns( tfe, knots=t.kn ) + age + factor( sex ),
+ family=poisreg, data=Lung.s, eps=10^-8, maxit=25 )
+ )

user system elapsed
0.271 0.459 0.240
```

Who needs the Cox-model anyway? (RXCox)

73/ 333

Mayo Clinic lung cancer 60 year old woman



Who needs the Cox-model anyway? (RXCox)

69/ 333

Example: Mayo Clinic lung cancer V

```
> library( mgcv )
> system.time(
+ mLs.pois.ps <- gam( cbind(lex.Xst=="Dead",lex.dur) ~
+ s( tfe ) + age + factor( sex ),
+ family=poisreg, data=Lung.s, eps=10^-8, maxit=25 )
+ )

user system elapsed
1.697 2.756 1.270

> ests <-
+ rbind( ci.exp(mL.cox),
+ ci.exp(mLs.pois.fc,subset=c("age","sex")),
+ ci.exp(mLs.pois.sp,subset=c("age","sex")),
+ ci.exp(mLs.pois.ps,subset=c("age","sex")))
> cmp <- cbind( ests[c(1,3,5,7) ,],
+ ests[c(1,3,5,7)+1,] )
> rownames( cmp ) <- c("Cox","Poisson-factor","Poisson-spline","Poisson-Pspline")
> colnames( cmp )[c(1,4)] <- c("age","sex")
```

Who needs the Cox-model anyway? (RXCox)

74/ 333

Example: Mayo Clinic lung cancer I

```
> library( survival )
> library( Epi )
> Lung <- Lexis( exit = list( tfe=time ),
+ exit.status = factor(status,labels=c("Alive","Dead")),
+ data = lung )
```

NOTE: entry.status has been set to "Alive" for all.
NOTE: entry is assumed to be 0 on the tfe timescale.

```
> summary( Lung )
```

Transitions:

From	To	Alive	Dead	Records	Events	Risk time	Persons
Alive	Alive	63	165	228	165	69593	228

Who needs the Cox-model anyway? (RXCox)

70/ 333

Example: Mayo Clinic lung cancer VI

```
> round( cmp, 7 )

age 2.5% 97.5% sex 2.5% 97.5%
Cox 1.017158 0.9989388 1.035710 0.5989574 0.4313720 0.8316487
Poisson-factor 1.017158 0.9989388 1.035710 0.5989574 0.4313720 0.8316487
Poisson-spline 1.016189 0.9980321 1.034677 0.5998287 0.4319854 0.8328858
Poisson-Pspline 1.016419 0.9982554 1.034913 0.6031306 0.4345167 0.8371751
```

Who needs the Cox-model anyway? (RXCox)

75/ 333

Example: Mayo Clinic lung cancer II

```
> system.time(
+ mL.cox <- coxph( Surv( tfe, tfe+lex.dur, lex.Xst=="Dead" ) ~
+ age + factor( sex ),
+ method="breslow", data=Lung ) )

user system elapsed
0.029 0.013 0.023


> Lung.s <- splitLexis( Lung,
+ breaks=c(0,sort(unique(Lung$time))),
+ time.scale="tfe" )
> summary( Lung.s )

Transitions:
To
From Alive Dead Records Events Risk time Persons
Alive 19857 165 20022 165 69593 228

> subset( Lung.s, lex.id==96 )[,1:11] ; nlevels( factor( Lung.s$tfe ) )
```

Who needs the Cox-model anyway? (RXCox)

71/ 333



Who needs the Cox-model anyway? (RXCox)

76/ 333

Example: Mayo Clinic lung cancer III

```
lex.id tfe lex.dur lex.Cst lex.Xst inst time status age sex ph.ecog
9235 96 0 5 Alive Alive 12 30 2 72 1 2
9236 96 5 6 Alive Alive 12 30 2 72 1 2
9237 96 11 1 Alive Alive 12 30 2 72 1 2
9238 96 12 1 Alive Alive 12 30 2 72 1 2
9239 96 13 2 Alive Alive 12 30 2 72 1 2
9240 96 15 11 Alive Alive 12 30 2 72 1 2
9241 96 26 4 Alive Dead 12 30 2 72 1 2


[1] 186

> system.time(
+ mLs.pois.fc <- glm( cbind(lex.Xst=="Dead", lex.dur) ~
+ - 1 + factor( tfe ) + age + factor( sex ),
+ family=poisreg, data=Lung.s, eps=10^-8, maxit=25 )
+ )

user system elapsed
12.529 18.409 8.871
```


Who needs the Cox-model anyway? (RXCox)

72/ 333



Who needs the Cox-model anyway? (RXCox)

76/ 333



Who needs the Cox-model anyway? (RNCox) 76/ 333

Follow-up and rates

- ▶ In follow-up studies we estimate rates from:
 - ▶ D — events, deaths
 - ▶ Y — person-years
 - ▶ $\hat{\lambda} = D/Y$ rates
 - ▶ ... empirical counterpart of intensity — **estimate**
- ▶ Rates differ between persons.
- ▶ Rates differ **within** persons:
 - ▶ By age
 - ▶ By calendar time
 - ▶ By disease duration
 - ▶ ...
- ▶ Multiple timescales.
- ▶ Multiple states (little boxes — later)

Follow-up data (time-split) 80/ 333

Deriving the survival function

```
> mLS.pois.sp <- glm( cbind(lex.Xst=="Dead",lex.dur) ~
+ Ns( tfe, knots=t.kn ) + age + factor( sex ),
+ family=poisreg, data=Lung.s, eps=10^-8, maxit=25 )

> CM <- cbind( 1, Ns( seq(10,1000,10)-5, knots=t.kn ), 60, 1 )
> lambda <- ci.exp( mLS.pois.sp, ctr.mat=CM )
> survP <- ci.surv( mLS.pois.sp, ctr.mat=CM, int1=10 )[,-4]
```

Code and output for the entire example available in <http://bendixcarstensen.com/AdvCoh/WNtCMA/>


Who needs the Cox-model anyway? (RNCox) 77/ 333

Examples: stratification by age

If follow-up is rather short, age at entry is OK for age-stratification.

If follow-up is long, use stratification by categories of **current age**, both for:

No. of events, D , and Risk time, Y .



— assuming a constant rate λ throughout.

Follow-up data (time-split) 81/ 333

What the Cox-model really is

Taking the life-table approach *ad absurdum* by:

- ▶ dividing time very finely and
 - ▶ modeling one covariate, the time-scale, with one parameter per distinct value.
 - ▶ the **model** for the time scale is really with exchangeable time-intervals.
- ⇒ difficult to access the baseline hazard (which looks terrible)
- ⇒ uninitiated tempted to show survival curves where irrelevant

Code and output for the entire example available in <http://bendixcarstensen.com/AdvCoh/WNtCMA/>

Who needs the Cox-model anyway? (RNCox) 78/ 333

Representation of follow-up data

A cohort or follow-up study records:

Events and Risk time.

The outcome is thus **bivariate**: (d, y)

Follow-up **data** for each individual must therefore have (at least) three variables:

Date of entry	entry	date variable
Date of exit	exit	date variable
Status at exit	fail	indicator (0/1)

Specific for each **type** of outcome.

Follow-up data (time-split) 82/ 333

Models of this world

- ▶ Replace the α_{it} s by a parametric function $f(t)$ with a limited number of parameters, for example:
 - ▶ Piecewise constant
 - ▶ Splines (linear, quadratic or cubic)
 - ▶ Fractional polynomials
- ▶ the two latter brings model into "this world":
 - ▶ smoothly varying rates
 - ▶ parametric closed form representation of baseline hazard
 - ▶ finite no. of parameters
- ▶ Makes it really easy to use rates directly in calculations of
 - ▶ expected residual life time
 - ▶ state occupancy probabilities in multistate models
 - ▶ ...

Who needs the Cox-model anyway? (RNCox) 79/ 333

Probability	log-Likelihood
$P(d \text{ at } t_x \text{entry } t_0)$	$d \log(\lambda) - \lambda y$
$= P(\text{surv } t_0 \rightarrow t_1 \text{entry } t_0)$	$= 0 \log(\lambda_1) - \lambda_1 y_1$
$\times P(\text{surv } t_1 \rightarrow t_2 \text{entry } t_1)$	$+ 0 \log(\lambda_2) - \lambda_2 y_2$
$\times P(d \text{ at } t_x \text{entry } t_2)$	$+ d \log(\lambda_3) - \lambda_3 y_3$

— allows different rates (λ_i) in each interval

Follow-up data (time-split) 83/ 333

Follow-up data

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020> time-split

Dividing time into bands:

If we want to compute D and Y in intervals on some timescale we must decide on:

Origin: The date where the time scale is 0:

- ▶ Age — 0 at date of birth
- ▶ Disease duration — 0 at date of diagnosis
- ▶ Occupation exposure — 0 at date of hire

Intervals: How should it be subdivided:

- ▶ 1-year classes? 5-year classes?
- ▶ Equal length?

Aim: Separate rate in each interval

Follow-up data (time-split) 84/ 333

Example: cohort with 3 persons:

Id	Bdate	Entry	Exit	St
1	14/07/1952	04/08/1965	27/06/1997	1
2	01/04/1954	08/09/1972	23/05/1995	0
3	10/06/1987	23/12/1991	24/07/1998	1

- ▶ Age bands: 10-years intervals of current age.
- ▶ Split Y for every subject accordingly
- ▶ Treat each segment as a separate unit of observation.
- ▶ Keep track of exit status in each interval.

Follow-up data (time-split)

85 / 333

Follow-up on several timescales

- ▶ The risk-time is the same on all timescales
- ▶ Only need the entry point on each time scale:
 - ▶ Age at entry.
 - ▶ Date of entry.
 - ▶ Time since treatment at entry.
 - if time of treatment is the entry, this is 0 for all.

- ▶ **Response variable** in analysis of rates:

(d, y) (event, duration)

- ▶ **Covariates** in analysis of rates:

- ▶ timescales
- ▶ other (fixed) measurements

- ▶ ... do not confuse **duration** and **timescale** !

Follow-up data (time-split)

90 / 333

Splitting the follow up

	subj. 1	subj. 2	subj. 3
Age at Entry:	13.06	18.44	4.54
Age at eXit:	44.95	41.14	11.12
Status at exit:	Dead	Alive	Dead
<hr/>			
Y	31.89	22.70	6.58
D	1	0	1

Follow-up data (time-split)

86 / 333

Follow-up data in Epi — Lexis objects

> thoro[1:6,1:8]

```
id sex birthdat contrast injecdat volume exitdat exitstat
1 1 2 1916.609 1 1938.791 22 1976.787 1
2 2 2 1927.843 1 1943.906 80 1966.030 1
3 3 1 1902.778 1 1935.629 10 1959.719 1
4 4 1 1918.359 1 1936.396 10 1977.307 1
5 5 1 1902.931 1 1937.387 10 1945.387 1
6 6 2 1903.714 1 1937.316 20 1944.738 1
```

Timescales of interest:

- ▶ Age
- ▶ Calendar time
- ▶ Time since injection

Follow-up data (time-split)

91 / 333

	subj. 1		subj. 2		subj. 3		Σ	
Age	Y	D	Y	D	Y	D	Y	D
0-	0.00	0	0.00	0	5.46	0	5.46	0
10-	6.94	0	1.56	0	1.12	1	8.62	1
20-	10.00	0	10.00	0	0.00	0	20.00	0
30-	10.00	0	10.00	0	0.00	0	20.00	0
40-	4.95	1	1.14	0	0.00	0	6.09	1
Σ	31.89	1	22.70	0	6.58	1	60.17	2

Follow-up data (time-split)

87 / 333

Definition of Lexis object

```
thL <- Lexis( entry = list( age = injecdat-birthdat,
                           per = injecdat,
                           tfi = 0 ),
              exit = list( per = exitdat ),
              exit.status = as.numeric(exitstat==1),
              data = thoro )
```

entry is defined on **three** timescales,

but **exit** is only needed on **one** timescale:

Follow-up time is the same on all timescales:

$exitdat - injecdat$

One element of **entry** and **exit** must have same name (**per**).

Follow-up data (time-split)

92 / 333

Splitting the follow-up

id	Bdate	Entry	Exit	St	risk	int
1	14/07/1952	03/08/1965	14/07/1972	0	6.9432	10
1	14/07/1952	14/07/1972	14/07/1982	0	10.0000	20
1	14/07/1952	14/07/1982	14/07/1992	0	10.0000	30
1	14/07/1952	14/07/1992	27/06/1997	1	4.9528	40
2	01/04/1954	08/09/1972	01/04/1974	0	1.5606	10
2	01/04/1954	01/04/1974	31/03/1984	0	10.0000	20
2	01/04/1954	31/03/1984	01/04/1994	0	10.0000	30
2	01/04/1954	01/04/1994	23/05/1995	0	1.1417	40
3	10/06/1987	23/12/1991	09/06/1997	0	5.4634	0
3	10/06/1987	09/06/1997	24/07/1998	1	1.1211	10

Keeping track of calendar time too?

Follow-up data (time-split)

88 / 333

The looks of a Lexis object

```
> thL[1:4,1:9]
  age per tfi lex.dur lex.Cst lex.Xst lex.id
1 22.18 1938.79 0 37.99 0 1 1
2 49.54 1945.77 0 18.59 0 1 2
3 68.20 1955.18 0 1.40 0 1 3
4 20.80 1957.61 0 34.52 0 0 4
...
```

```
> summary( thL )
Transitions:
To
From 0 1 Records: Events: Risk time: Persons:
0 504 1964 2468 1964 51934.08 2468
```

Follow-up data (time-split)


93 / 333

Timescales

- ▶ A timescale is a variable that varies **deterministically** *within* each person during follow-up:
 - ▶ Age
 - ▶ Calendar time
 - ▶ Time since treatment
 - ▶ Time since relapse
- ▶ All timescales advance at the same pace (1 year per year ...)
- ▶ Note: Cumulative exposure is **not** a timescale.

Follow-up data (time-split)


89 / 333



> plot(thL, lwd=3)


Follow-up data (time-split)

94 / 333



Likelihood for a constant rate

- ▶ This setup is for a situation where it is assumed that rates are constant in each of the intervals.
- ▶ Each observation in the dataset contributes a term to the likelihood.
- ▶ Each term looks like a contribution from a Poisson variate (albeit with values only 0 or 1)
- ▶ Rates can vary along several timescales simultaneously.
- ▶ Models can include fixed covariates, as well as the timescales (the left end-points of the intervals) as **quantitative** variables.
- ▶ The latter is where we will need splines.



The Poisson likelihood for split data

- ▶ Split records (one per person-interval (p, i)):

$$\sum_{p,i} (d_{pi} \log(\lambda) - \lambda y_{pi}) = D \log(\lambda) - \lambda Y$$

- ▶ Assuming that the death indicator ($d_{pi} \in \{0, 1\}$) is Poisson, a model with with offset $\log(y_{pi})$ will give the same result.
- ▶ If we assume that rates are constant we get the simple expression with (D, Y)
- ▶ ... but the split data allows models that assume different rates for different (d_{pi}, y_{pi}) , so rates can vary **within** a person's follow-up.

Splitting follow-up time

```

> spl1 <- splitLexis( thL, breaks=seq(0,100,20),
+                   time.scale="age" )
> round(spl1,1)

```

age	per	tfi	lex.dur	lex.Cst	lex.Xst	id	sex	birthdat	contrast	injecdat	vo
1	22.2	1938.8	0.0	17.8	0	0	1	2	1916.6	1	1938.8
2	40.0	1956.6	17.8	20.0	0	0	1	2	1916.6	1	1938.8
3	60.0	1976.6	37.8	0.2	0	1	1	2	1916.6	1	1938.8
4	49.5	1945.8	0.0	10.5	0	0	640	2	1896.2	1	1945.8
5	60.0	1956.2	10.5	8.1	0	1	640	2	1896.2	1	1945.8
6	68.2	1955.2	0.0	1.4	0	1	3425	1	1887.0	2	1955.2
7	20.8	1957.6	0.0	19.2	0	0	4017	2	1936.8	2	1957.6
8	40.0	1976.8	19.2	15.3	0	0	4017	2	1936.8	2	1957.6

Follow-up data (time-split) 96/ 333

Where is (d_{pi}, y_{pi}) in the split data?

```

> spl1 <- splitLexis( thL , breaks=seq(0,100,20) , time.scale="age" )
> spl2 <- splitLexis( spl1, breaks=c(0,1,5,20,100), time.scale="tfi" )
> options( digits=5 )
> spl2[1:10,1:11]

```

lex.id	age	per	tfi	lex.dur	lex.Cst	lex.Xst	id	sex	birthdat	contrast	vo
1	1	22.182	1938.8	0.000	1.00000	0	0	1	2	1916.6	1
2	1	23.182	1939.8	1.000	4.00000	0	0	1	2	1916.6	1
3	1	27.182	1943.8	5.000	12.81793	0	0	1	2	1916.6	1
4	1	40.000	1956.6	17.818	2.18207	0	0	1	2	1916.6	1
5	1	42.182	1958.8	20.000	17.81793	0	0	1	2	1916.6	1
6	1	60.000	1976.6	37.818	0.17796	0	1	1	2	1916.6	1
7	2	16.063	1943.9	0.000	1.00000	0	0	2	2	1927.8	1
8	2	17.063	1944.9	1.000	2.93703	0	0	2	2	1927.8	1
9	2	20.000	1947.8	3.937	1.06297	0	0	2	2	1927.8	1
10	2	21.063	1948.9	5.000	15.00000	0	0	2	2	1927.8	1

— and what are covariates for the rates?

Split on another timescale

```

> spl2 <- splitLexis( spl1, time.scale="tfi",
+                   breaks=c(0,1,5,20,100) )
> round( spl2, 1 )

```

lex.id	age	per	tfi	lex.dur	lex.Cst	lex.Xst	id	sex	birthdat	contrast	inje
1	1	22.2	1938.8	0.0	1.0	0	0	1	2	1916.6	1
2	1	23.2	1939.8	1.0	4.0	0	0	1	2	1916.6	1
3	1	27.2	1943.8	5.0	12.8	0	0	1	2	1916.6	1
4	1	40.0	1956.6	17.8	2.2	0	0	1	2	1916.6	1
5	1	42.2	1958.8	20.0	17.8	0	0	1	2	1916.6	1
6	1	60.0	1976.6	37.8	0.2	0	1	1	2	1916.6	1
7	2	49.5	1945.8	0.0	1.0	0	0	640	2	1896.2	1
8	2	50.5	1946.8	1.0	4.0	0	0	640	2	1896.2	1
9	2	54.5	1950.8	5.0	5.5	0	0	640	2	1896.2	1
10	2	60.0	1956.2	10.5	8.1	0	1	640	2	1896.2	1
11	3	68.2	1955.2	0.0	1.0	0	1	3425	1	1887.0	2
12	3	69.2	1956.2	1.0	0.4	0	1	3425	1	1887.0	2
13	4	20.8	1957.6	0.0	1.0	0	0	4017	2	1936.8	2
14	4	21.8	1958.6	1.0	4.0	0	0	4017	2	1936.8	2
15	4	25.8	1962.6	5.0	14.2	0	0	4017	2	1936.8	2
16	4	40.0	1976.8	19.2	0.8	0	0	4017	2	1936.8	2
17	4	40.8	1977.6	20.0	14.5	0	0	4017	2	1936.8	2

Follow-up data (time-split) 98/ 333

Where is (d_{pi}, y_{pi}) in the split data?


```

> library( popEpi )
> spl1 <- splitMulti( thL , age=seq(0,100,20) )
> spl2 <- splitMulti( spl1, tfi=c(0,1,5,20,100) )
> options( digits=5 )
> spl2[1:10,1:11]

```

lex.id	age	per	tfi	lex.dur	lex.Cst	lex.Xst	id	sex	birthdat	contrast	vo
1:	1	22.182	1938.8	0.000	1.00000	0	0	1	2	1916.6	1
2:	1	23.182	1939.8	1.000	4.00000	0	0	1	2	1916.6	1
3:	1	27.182	1943.8	5.000	12.81793	0	0	1	2	1916.6	1
4:	1	40.000	1956.6	17.818	2.18207	0	0	1	2	1916.6	1
5:	1	42.182	1958.8	20.000	17.81793	0	0	1	2	1916.6	1
6:	1	60.000	1976.6	37.818	0.17796	0	1	1	2	1916.6	1
7:	2	16.063	1943.9	0.000	1.00000	0	0	2	2	1927.8	1
8:	2	17.063	1944.9	1.000	2.93703	0	0	2	2	1927.8	1
9:	2	20.000	1947.8	3.937	1.06297	0	0	2	2	1927.8	1
10:	2	21.063	1948.9	5.000	15.00000	0	0	2	2	1927.8	1

— not the printing: it's a `data.table`



Analysis of results

- ▶ d_{pi} — events in the variable: `lex.Xst`:
In the model as response: `lex.Xst==1`
- ▶ y_{pi} — risk time: `lex.dur` (duration):
In the model as offset $\log(y)$, $\log(\text{lex.dur})$.
- ▶ Covariates are:
 - ▶ timescales (age, period, time in study)
 - ▶ other variables for this person (constant or *assumed* constant in each interval).
- ▶ Model rates using the covariates in `glm`:
— no difference between time-scales and other covariates.

Fitting a simple model

```
> stat.table( contrast,
+             list( D = sum( lex.Xst ),
+                 Y = sum( lex.dur ),
+                 Rate = ratio( lex.Xst, lex.dur, 100 ) ),
+             margin = TRUE,
+             data = spl2 )
```

contrast	D	Y	Rate
1	928.00	20094.74	4.62
2	1036.00	31822.24	3.26
Total	1964.00	51916.98	3.78

Follow-up data (time-split)

104/ 333

Practical set-up

- ▶ Tabulate only events (as obtained from the cancer registry) by age and period.
- ▶ 100 age-classes.
- ▶ 65 periods (single calendar years).
- ▶ 6500 aggregate records of D .
- ▶ Estimate the population follow-up based on census data from Statistics Denmark (Y_{pop}).
... or get it from the human mortality database.
- ▶ If disease is common: tabulate follow-up **after** diagnosis (Y_{dis}), and subtract from population follow-up.
- ▶ Analyze (D , Y) by Poisson model.

Models for tabulated data (tab-mod)

108/ 333

Fitting a simple model

contrast	D	Y	Rate
1	928.00	20094.74	4.62
2	1036.00	31822.24	3.26


```
> m0 <- glm( (lex.Xst==1) ~ factor(contrast) - 1,
+           offset = log(lex.dur/100),
+           family = poisson,
+           data = spl2 )
> round( ci.exp( m0 ), 2 )
```

```
exp(Est.) 2.5% 97.5%
factor(contrast)1 4.62 4.33 4.93
factor(contrast)2 3.26 3.06 3.46
```

Follow-up data (time-split)

105/ 333

Lexis diagram ¹



Disease registers record events.

Official statistics collect population data.

¹ Named after the German statistician and economist William Lexis (1837–1914), who devised this diagram in the book "Einführung in die Theorie der Bevölkerungsstatistik" (Karl J. Trübner, Strassburg, 1875).

Models for tabulated data (tab-mod)

109/ 333

Models for tabulated data


Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

tab-mod

Lexis diagram



Registration of:

cases (D)

risk time,
person-years (Y)

in subsets of the Lexis
diagram.

Models for tabulated data (tab-mod)

110/ 333

Conceptual set-up


Follow-up of the entire (male) population from 1943–2006 w.r.t. occurrence of testis cancer:

- ▶ Split follow-up time for all about 4 mil. men in 1-year classes by age and calendar time (y).
- ▶ Allocate testis cancer event ($d = 0, 1$) to each.
- ▶ Analyze all 200,000,000 records by a Poisson model.

Models for tabulated data (tab-mod)

106/ 333

Lexis diagram



Registration of:

cases (D)

risk time,
person-years (Y)

in subsets of the Lexis
diagram.

Rates available in each
subset.

Models for tabulated data (tab-mod)

111/ 333

Realistic set-up

- ▶ Tabulate the follow-up time and events by age and period.
- ▶ 100 age-classes.
- ▶ 65 periods (single calendar years).
- ▶ 6500 aggregate records of (D , Y).
- ▶ Analyze by a Poisson model.

Models for tabulated data (tab-mod)

107/ 333

Register data

Classification of **cases** (D_{ap}) by age at diagnosis and date of diagnosis, and **population** (Y_{ap}) by age at risk and date at risk, in compartments of the Lexis diagram, e.g.:

```
> fCtable( xtable( cbind(D,Y) ~ A + P, data=ts ), col.vars=3:2, w=8 )
```

A	P	D				Y			
		1943	1948	1953	1958	1943	1948	1953	1958
15		2	3	4	1	773,812	744,217	794,123	972,853
20		7	7	17	8	813,022	744,706	721,810	770,859
25		28	23	26	35	790,501	781,827	722,968	698,612
30		28	43	49	51	799,293	774,542	769,298	711,596
35		36	42	39	44	769,356	782,893	760,213	760,452
40		24	32	46	53	694,073	754,322	768,471	749,912

Models for tabulated data (tab-mod)

112/ 333

In analysis format:

```
> ts
      A   P   D   Y
1  15 1943  2  773812
2  20 1943  7  813022
3  25 1943 28  790501
4  30 1943 28  799293
5  35 1943 36  769356
6  40 1943 24  694073
7  15 1948  3  744217
8  20 1948  7  744706
9  25 1948 23  781827
10 30 1948 43  774542
11 35 1948 42  782893
12 40 1948 32  754322
13 15 1953  4  794123
14 20 1953 17  721810
15 25 1953 26  722968
16 30 1953 49  769298
17 35 1953 39  760213
18 40 1953 46  684711
19 15 1958  1  729822
```

Models for tabulated data (tab-mod)

113 / 333

Linear combinations of the parameters can be computed using the `ctr.mat` option:

```
> CM <- rbind( '1943 vs. 1953' = c( 0,-1, 0),
+             '1948 vs. 1953' = c( 1,-1, 0),
+             'Ref. (1953)' = c( 0, 0, 0),
+             '1958 vs. 1953' = c( 0,-1, 1) )
> round( ci.exp( m0, subset="P", ctr.mat=CM ), 3 )
              exp(Est.)  2.5% 97.5%
1943 vs. 1953    0.882 0.543 0.857
1948 vs. 1953    0.813 0.655 1.010
Ref. (1953)      1.000 1.000 1.000
1958 vs. 1953    1.087 0.887 1.332
```

Models for tabulated data (tab-mod)

118 / 333

Tabulated data

Once data are in tabular form, models are restricted:

- ▶ Rates must be assumed constant in each cell of the table / subset of the Lexis diagram.
- ▶ With large cells (5 × 5 years) it is customary to put a separate parameter on each cell or on each levels of classifying factors.
- ▶ Output from the model will be rates and rate-ratios.
- ▶ Since we use multiplicative Poisson, usually the log rates and the log-RR are reported

Models for tabulated data (tab-mod)

114 / 333

Age-Period and Age-Cohort models

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:

Age-Period-Cohort models

— and some cousins

European Doctoral School of Demography,

Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

AP-AC

Simple age-period model for the testis cancer rates:

```
> m0 <- glm( D ~ factor(A) + factor(P) + offset(log(Y/10^5)),
+          family=poisson, data=ts )
> summary( m0 )

Call:
glm(formula = D ~ factor(A) + factor(P) + offset(log(Y/10^5)),
    family = poisson, data = ts)

Deviance Residuals:
    Min       1Q   Median       3Q      Max
-1.5991  -0.6974   0.1284   0.6671   1.8904

Coefficients:
              Estimate Std. Error z value Pr(>|z|)
(Intercept)  -1.4758    0.3267  -4.517 6.26e-06
factor(A)20    1.4539    0.3545   4.101 4.11e-05
factor(A)25    2.5321    0.3301   7.671 1.71e-14
factor(A)30    2.9327    0.3254   9.013 < 2e-16
factor(A)35    2.8613    0.3259   8.779 < 2e-16
factor(A)40    2.8521    0.3263   8.741 < 2e-16
factor(P)1948  0.1753    0.1211   1.447 0.14778
factor(P)1953  0.3822    0.1163   3.286 0.00102
```

Models for tabulated data (tab-mod)

115 / 333

Register data — rates

Rates in “tiles” of the Lexis diagram:

$$\lambda(a, p) = D_{ap} / Y_{ap}$$

Descriptive epidemiology based on disease registers:

How do the rates vary by age and time:

- ▶ Age-specific rates for a given period.
- ▶ Age-standardized rates as a function of calendar time. (Weighted averages of the age-specific rates).

Age-Period and Age-Cohort models (AP-AC)

119 / 333

`ci.exp()` from the Epi package extracts coefficients and computes confidence intervals:

```
> round( ci.exp( m0 ), 2 )
              exp(Est.)  2.5% 97.5%
(Intercept)    0.23 0.12 0.43
factor(A)20    4.28 2.14 8.57
factor(A)25   12.58 6.59 24.02
factor(A)30   18.78 9.92 35.53
factor(A)35   17.49 9.23 33.12
factor(A)40   17.32 9.14 32.84
factor(P)1948  1.19 0.94 1.51
factor(P)1953  1.47 1.17 1.84
factor(P)1958  1.59 1.27 2.00
```

... what do these parameters mean?

Models for tabulated data (tab-mod)

116 / 333

Subsets of parameter estimates accessed via a character string that is `grep`-ed to the names.


```
> round( ci.exp( m0, subset="P", pval=TRUE ), 3 )
              exp(Est.)  2.5% 97.5%   P
factor(P)1948  1.192 0.940 1.511 0.148
factor(P)1953  1.466 1.167 1.841 0.001
factor(P)1958  1.593 1.272 1.996 0.000

> round( ci.lin( m0, subset="P" ), 3 )
              Estimate StdErr      z      P  2.5% 97.5%
factor(P)1948  0.175  0.121 1.447 0.148 -0.062 0.413
factor(P)1953  0.382  0.116 3.286 0.001  0.154 0.610
factor(P)1958  0.466  0.115 4.052 0.000  0.241 0.691
```

Models for tabulated data (tab-mod)

117 / 333

“Synthetic” cohorts




Events and risk time in cells along the diagonals are among persons with roughly same date of birth.

Successively overlapping 10-year periods.

Age-Period and Age-Cohort models (AP-AC)

120 / 333

Lexis diagram: data



Testis cancer cases in Denmark.

Male person-years in Denmark.

Age-Period and Age-Cohort models (AP-AC)

121 / 333

Data matrix: Testis cancer cases

Number of cases

Age	Date of diagnosis (year - 1900)									
	48-52	53-57	58-62	63-67	68-72	73-77	78-82	83-87	88-92	
15-19	7	13	13	15	33	35	37	49	51	
20-24	31	46	49	55	85	110	140	151	150	
25-29	62	63	82	87	103	153	201	214	268	
30-34	66	82	88	103	124	164	207	209	258	
35-39	56	56	67	99	124	142	152	188	209	
40-44	47	65	64	67	85	103	119	121	155	
45-49	30	37	54	45	64	63	66	92	86	
50-54	28	22	27	46	36	50	49	61	64	
55-59	14	16	25	26	29	28	43	42	34	

```
> ts <- transform( subset( testisDK, A>14 & A<60 ),
+                 A = floor( A /5)*5 +2.5,
+                 P = floor((P-1943)/5)*5+1943+2.5 )
> ts$C <- ts$P - ts$A
> trate <- xtabs( D ~ A + P, data = ts ) /
+           xtabs( Y ~ A + P, data = ts ) * 100000
> #
> #
> #
> trate[1:5,1:6]
```

A	P	1945.5	1950.5	1955.5	1960.5	1965.5	1970.5
17.5	1.2923036	0.9405857	1.6370257	1.3362759	1.4264867	3.4340862	3.4340862
22.5	3.6899378	4.1627194	6.3728682	6.3565492	5.7274822	8.0657826	8.0657826
27.5	6.9576174	7.9301414	8.7140826	11.7375624	11.3753792	10.6996275	10.6996275
32.5	7.0061961	8.5211703	10.6590661	12.3665762	14.7122260	16.1068525	16.1068525
37.5	6.8888785	7.1529555	7.3663549	8.8105514	13.9126492	17.6571019	17.6571019

Data matrix: Male risk time

1000 person-years

Age	Date of diagnosis (year - 1900)									
	48-52	53-57	58-62	63-67	68-72	73-77	78-82	83-87	88-92	
15-19	744.2	794.1	972.9	1051.5	961.0	952.5	1011.1	1005.0	929.8	
20-24	744.7	721.8	770.9	960.3	1053.8	967.5	953.0	1019.7	1017.3	
25-29	781.8	723.0	698.6	764.8	962.7	1056.1	960.9	956.2	1031.6	
30-34	774.5	769.3	711.6	700.1	769.9	960.4	1045.3	955.0	957.1	
35-39	782.9	760.2	760.5	711.6	702.3	767.5	951.9	1035.7	948.6	
40-44	754.3	768.5	749.9	756.5	709.8	696.5	757.8	940.3	1023.7	
45-49	676.7	737.9	753.5	738.1	746.4	698.2	682.4	743.1	923.4	
50-54	600.3	653.9	715.4	732.7	718.3	724.2	675.5	660.8	721.1	
55-59	512.8	571.1	622.5	680.8	698.2	683.8	686.4	640.9	627.7	


```
> par( mfrow=c(2,2) )
> rateplot( trate, col=gray(2:15/18), lwd=3, ann=TRUE )
> wh = c("ap", "ac", "pa", "ca")
> for( ptp in wh ) {
+   pdf( paste("./AP-AC-", ptp, ".pdf", sep="" ), height=6, width=8 )
+   par( mar=c(3,3,1,1), mgp=c(3,1,0)/1.6, bty="n", las=1 )
+   rateplot( trate, which=ptp,
+            col=gray(2:15/18), lwd=3, ann=TRUE, a.lim=c(15,60) )
+   dev.off()
+ }
```

Data matrix: Empirical rates

Rate per 1000,000 person-years

Age	Date of diagnosis (year - 1900)									
	48-52	53-57	58-62	63-67	68-72	73-77	78-82	83-87	88-92	
15-19	9.4	16.4	13.4	14.3	34.3	36.7	36.6	48.8	54.8	
20-24	41.6	63.7	63.6	57.3	80.7	113.7	146.9	148.1	147.4	
25-29	79.3	87.1	117.4	113.8	107.0	144.9	209.2	223.8	259.8	
30-34	85.2	106.6	123.7	147.1	161.1	170.8	198.0	218.8	269.6	
35-39	71.5	73.7	88.1	139.1	176.6	185.0	159.7	181.5	220.3	
40-44	62.3	84.6	85.3	88.6	119.8	147.9	157.0	128.7	151.4	
45-49	44.3	50.1	71.7	61.0	85.7	90.2	96.7	123.8	93.1	
50-54	46.6	33.6	37.7	62.8	50.1	69.0	72.5	92.3	88.7	
55-59	27.3	28.0	40.2	38.2	41.5	40.9	62.6	65.5	54.2	

which = "ap"




The classical plots

Given a table of rates classified by age and period, we can do 4 "classical" plots:

- ▶ Rates versus age at diagnosis (period):
 - rates in the same age-class connected.
- ▶ Rates versus age at diagnosis:
 - rates in the same birth-cohort connected.
- ▶ Rates versus date of diagnosis:
 - rates in the same age-class connected.
- ▶ Rates versus date of date of birth:
 - rates in the same age-class connected.

These plots can be produced by the R-function `rateplot`.


which = "ac"




```
> library( Epi )
> data( testisDK )
> head( testisDK )
```

A	P	D	Y
1	0	1943	1 39649.50
2	1	1943	1 36942.83
3	2	1943	0 34588.33
4	3	1943	1 33267.00
5	4	1943	0 32614.00
6	5	1943	0 32020.33

which = "pa"



which = "ca"



Estimates with confidence intervals

```
> par( mfrow=c(1,2), mar=c(3,3,1,1), mgp=c(3,1,0)/1.6, bty="n", las=1 )
> matshade( seq(17.5,57.5,5), ci.exp(ap,subset="A"), plot=TRUE,
+ log="y", lwd=2, ylim=c(1,20), xlab="Age",
+ ylab="Testis cancer rate per 100,000 PY (1970)" )
> matshade( seq(1945.5,1995.5,5),
+ rbind( ci.exp(ap,subset="P")[1:5,], 1,
+ ci.exp(ap,subset="P")[6:10,] ), plot=TRUE,
+ log="y", lwd=2, ylim=c(1,20)/5,
+ xlab="Date of follow-up", ylab="Rate ratio" )
> abline( h = 1 )
> points( 1970.5, 1, pch=16 )
```

Age-Period model


Rates are proportional between periods:

$$\lambda(a, p) = a_a \times b_p \quad \text{or} \quad \log[\lambda(a, p)] = \alpha_a + \beta_p$$

Choose p_0 as reference period, where $\beta_{p_0} = 0$

$$\log[\lambda(a, p_0)] = \alpha_a + \beta_{p_0} = \alpha_a$$

Estimates from Age-Period model



Fitting the A-P model in R I

Reference period is the 5th period (1970.5 ~ 1968-72):

```
> ap <- glm( D ~ factor(A) - 1 + relevel( factor(P), "1970.5" ) +
+ offset( log(Y/10^5) ),
+ family=poisson, data=ts )
> summary( ap )
```

Call:
glm(formula = D ~ factor(A) - 1 + relevel(factor(P), "1970.5") +
offset(log(Y/10^5)), family = poisson, data = ts)

Deviance Residuals:
Min 1Q Median 3Q Max
-3.1850 -0.9465 -0.1683 0.5767 3.8610

Coefficients:
Estimate Std. Error z value Pr(>|z|)

Age-cohort model

Rates are proportional between cohorts:

$$\lambda(a, c) = a_a \times c_c \quad \text{or} \quad \log[\lambda(a, p)] = \alpha_a + \gamma_c$$

Choose c_0 as reference cohort, where $\gamma_{c_0} = 0$

$$\log[\lambda(a, c_0)] = \alpha_a + \gamma_{c_0} = \alpha_a$$

Fitting the A-P model in R II

```
factor(A)17.5 1.06715 0.06791 15.715 < 2e-16
factor(A)22.5 2.20732 0.04837 45.630 < 2e-16
factor(A)27.5 2.65463 0.04465 59.449 < 2e-16
factor(A)32.5 2.77057 0.04458 62.142 < 2e-16
factor(A)37.5 2.63081 0.04606 57.122 < 2e-16
factor(A)42.5 2.36224 0.04878 48.422 < 2e-16
factor(A)47.5 2.01945 0.05341 37.811 < 2e-16
factor(A)52.5 1.73119 0.05957 29.062 < 2e-16
factor(A)57.5 1.45070 0.06748 21.498 < 2e-16
relevel(factor(P), "1970.5")1945.5 -0.75072 0.07011 -10.708 < 2e-16
relevel(factor(P), "1970.5")1950.5 -0.59740 0.06633 -9.006 < 2e-16
relevel(factor(P), "1970.5")1955.5 -0.43562 0.06299 -6.916 4.65e-12
relevel(factor(P), "1970.5")1960.5 -0.27952 0.05999 -4.659 3.18e-06
relevel(factor(P), "1970.5")1965.5 -0.16989 0.05751 -2.954 0.00313
relevel(factor(P), "1970.5")1975.5 0.16037 0.05143 3.118 0.00182
relevel(factor(P), "1970.5")1980.5 0.30022 0.04953 6.061 1.35e-09
relevel(factor(P), "1970.5")1985.5 0.37491 0.04853 7.726 1.11e-14
relevel(factor(P), "1970.5")1990.5 0.47047 0.04745 9.916 < 2e-16
relevel(factor(P), "1970.5")1995.5 0.54079 0.04862 11.123 < 2e-16
```

Fitting the A-C model in R I

Reference period is the 1933 cohort:

```
> ac <- glm( D ~ factor(A) - 1 + relevel( factor(C), "1933" ) +
+ offset( log(Y/10^5) ),
+ family=poisson, data=ts )
> summary( ac )
```

Call:
glm(formula = D ~ factor(A) - 1 + relevel(factor(C), "1933") +
offset(log(Y/10^5)), family = poisson, data = ts)

Deviance Residuals:
Min 1Q Median 3Q Max
-3.0796 -0.9538 -0.1620 0.5767 3.9525

Coefficients:
Estimate Std. Error z value Pr(>|z|)
factor(A)17.5 0.61513 0.07534 8.165 3.23e-16

Fitting the A-P model in R III

(Dispersion parameter for poisson family taken to be 1)

Null deviance: 29193.6 on 2430 degrees of freedom
Residual deviance: 2816.6 on 2411 degrees of freedom
AIC: 9005

Number of Fisher Scoring iterations: 5

Fitting the A-C model in R II

```
factor(A)22.5 1.89965 0.05342 35.558 < 2e-16
factor(A)27.5 2.46911 0.04842 50.990 < 2e-16
factor(A)32.5 2.70635 0.04695 57.639 < 2e-16
factor(A)37.5 2.71211 0.04758 57.006 < 2e-16
factor(A)42.5 2.58676 0.04993 51.803 < 2e-16
factor(A)47.5 2.36542 0.05459 43.327 < 2e-16
factor(A)52.5 2.18192 0.06098 35.782 < 2e-16
factor(A)57.5 2.01519 0.06939 29.041 < 2e-16
relevel(factor(C), "1933")1888 -1.77316 0.41400 -4.283 1.84e-05
relevel(factor(C), "1933")1893 -1.05641 0.19017 -5.555 2.77e-08
relevel(factor(C), "1933")1898 -0.79897 0.12600 -6.341 2.28e-10
relevel(factor(C), "1933")1903 -0.87599 0.10389 -8.432 < 2e-16
relevel(factor(C), "1933")1908 -0.76707 0.08352 -9.184 < 2e-16
relevel(factor(C), "1933")1913 -0.56290 0.07006 -8.035 9.36e-16
relevel(factor(C), "1933")1918 -0.56702 0.06683 -8.484 < 2e-16
relevel(factor(C), "1933")1923 -0.36836 0.06124 -6.015 1.79e-09
relevel(factor(C), "1933")1928 -0.18832 0.05903 -3.190 0.001421
relevel(factor(C), "1933")1938 0.08958 0.05439 1.647 0.099585
relevel(factor(C), "1933")1943 -0.03107 0.05443 -0.571 0.568091
```


Fitting the A-C model in R III

```
relevel(factor(C), "1933")1948 0.18088 0.05256 3.441 0.000579
relevel(factor(C), "1933")1953 0.42239 0.05309 7.956 1.77e-15
relevel(factor(C), "1933")1958 0.62544 0.05421 11.537 < 2e-16
relevel(factor(C), "1933")1963 0.75687 0.05727 13.215 < 2e-16
relevel(factor(C), "1933")1968 0.75183 0.06799 11.057 < 2e-16
relevel(factor(C), "1933")1973 0.87343 0.09373 9.318 < 2e-16
relevel(factor(C), "1933")1978 1.19601 0.17340 6.898 5.29e-12
```

(Dispersion parameter for poisson family taken to be 1)

```
Null deviance: 29193.6 on 2430 degrees of freedom
Residual deviance: 2767.8 on 2403 degrees of freedom
AIC: 8972.2
```

Number of Fisher Scoring iterations: 5

Recap of Monday — rates

- ▶ Rate, intensity: $\lambda(t) = P\{\text{event in } (t, t+h) | \text{alive at } t\} / h$
- ▶ Observed empirical rates (d, y) — possibly many per person.
- ▶ $\ell_{FU} = d \log(\lambda) - \lambda y$, obs: (d, y) , rate parameter: λ
- ▶ $\ell_{\text{Poisson}} = d \log(\lambda y) - \lambda y$, obs: d , mean parameter: $\mu = \lambda y$
- ▶ — use either to find m.l.e. of λ
- ▶ Poisson model is for $\log(\mu) = \log(\lambda) + \log(y)$ hence `offset=log(Y)` in `poisson`
- ▶ Better to use `glm(cbind(D,Y) ~ ..., family=poisreg)`
- ▶ Once rates are known, we can construct survival curves and derivatives of these.


Estimates with confidence intervals

```
> par(mfrow=c(1,2), mar=c(3,3,1,1), mgp=c(3,1,0)/1.6, bty="n", las=1)
> matshade(seq(17.5,57.5,5),
+ ci.exp(ac,subset="A"),
+ plot=TRUE,
+ log="y", lwd=2, ylim=c(1,20), xlab="Age",
+ ylab="Testis cancer rate per 100,000 PY (1933 cohort)")
> matshade(seq(1888,1978,5),
+ rbind(ci.exp(ac,subset="C")[1:9,], 1,
+ ci.exp(ac,subset="C")[10:18,]),
+ plot=TRUE,
+ log="y", lwd=2, ylim=c(1,20)/5,
+ xlab="Date of birth", ylab="Rate ratio")
> abline(h = 1)
> points(1933, 1, pch=16)
```

Recap Monday — models

- ▶ Empirical rate (d_t, y_t) relates to a **time** t
- ▶ Many for the same person — different times
- ▶ Not independent, but likelihood is a product
- ▶ One parameter per interval \Rightarrow exchangeable times
- ▶ Use the quantitative nature of t : \Rightarrow smooth continuous effects of time
- ▶ Predicted rates: `ci.pred(model, newdata=nd)`
- ▶ RR is the difference between two predictions:
- ▶ RR by period:
 - ▶ `ndx <- data.frame(P=1947:1980, A=47)`
 - ▶ `ndr <- data.frame(P=1970, A=47)`
 - ▶ `ci.exp(model, ctr.mat=list(ndx-ndr))`

Estimates from Age-Cohort model



Age-drift model

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

Monday exercise: Age-Period and Age-Cohort models

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

Linear effect of period:

$$\log[\lambda(a, p)] = \alpha_a + \beta_p = \alpha_a + \beta(p - p_0)$$

that is, $\beta_p = \beta(p - p_0)$.

Linear effect of cohort:

$$\log[\lambda(a, p)] = \tilde{\alpha}_a + \gamma_c = \tilde{\alpha}_a + \gamma(c - c_0)$$

that is, $\gamma_c = \gamma(c - c_0)$

Monday afternoon exercise

1. Load the `Epi` package.
2. Do the practicals 2.1 and 2.2, note where the data is.
3. I will be available the first approx. hour for direct questions that will be broadcast to the entire audience.
4. After that I will be on e-mail, until about 15.
5. Tuesday morning we will:
 - ▶ Recap the contents of today's lecture
 - ▶ Go over the solutions to today's exercises
6. Remember: Questions always welcome at any time, just switch on your microphone and peak.

Age and linear effect of period:

```
> apd <- glm(D ~ factor(A) - 1 + I(P-1970.5) +
+ offset(log(Y)),
+ family=poisson)
> summary(apd)
```

```
Call:
glm(formula = D ~ factor(A) - 1 + I(P - 1970.5) + offset(log(Y)), family = poisson)
```

```
Deviance Residuals:
    Min       1Q   Median       3Q      Max
-2.97593  -0.77091  0.02809  0.95914  2.93076
```

```
Coefficients:
            Estimate Std. Error z value Pr(>|z|)
factor(A)17.5 -3.58065     0.06306  -56.79  <2e-16
...
factor(A)57.5 -3.17579     0.06256  -50.77  <2e-16
I(P - 1970.5) 0.02653     0.00100   26.52  <2e-16
```

(Dispersion parameter for poisson family taken to be 1)

```
Null deviance: 89358.53 on 81 degrees of freedom
Residual deviance: 126.07 on 71 degrees of freedom
```

Age and linear effect of cohort:

```
> acd <- glm( D ~ factor( A ) - 1 + I( C - 1933 ) +
+           offset( log( Y ) ),
+           family = poisson )
> summary( acd )

Call:
glm(formula = D ~ factor(A) - 1 + I(C - 1933) + offset(log(Y)), family = poisson)

Deviance Residuals:
    Min       1Q   Median       3Q      Max
-2.97593 -0.77091  0.02809  0.95914  2.93076

Coefficients:
            Estimate Std. Error z value Pr(>|z|)
factor(A) 17.5 -4.11117   0.06760  -60.82 <2e-16
...
factor(A) 57.5 -2.64527   0.06423  -41.19 <2e-16
I(C - 1933)  0.02653   0.00100   26.52 <2e-16

(Dispersion parameter for poisson family taken to be 1)

Null deviance: 89358.53  on 81  degrees of freedom
Residual deviance: 126.07  on 71  degrees of freedom
```

Age-Period-Cohort model (APC-cat) 150 / 333

Fitting the model in R I

```
> library( Epi )
> data( testisDK )
> tc <- transform( subset( testisDK, A>14 & A<60 & P<1993 ),
+                 A = floor( A / 5 ) * 5 + 2.5,
+                 P = floor( ( P - 1943 ) / 5 ) * 5 + 1943 + 2.5 )
> tc <- aggregate( tc[, c( "D", "Y" ) ], tc[, c( "A", "P" ) ], FUN = sum )
> tc$C <- tc$P - tc$A
> str( tc )

'data.frame': 90 obs. of 5 variables:
 $ A: num  17.5 22.5 27.5 32.5 37.5 42.5 47.5 52.5 57.5 17.5 ...
 $ P: num  1946 1946 1946 1946 1946 ...
 $ D: num  10 30 55 56 53 35 29 16 6 7 ...
 $ Y: num  773812 813022 790500 799293 769356 ...
 $ C: num  1928 1923 1918 1913 1908 ...
```

Age-Period-Cohort model (APC-cat) 154 / 333

What goes on?

$$p = a + c \quad p_0 = a_0 + c_0$$

$$\alpha_a + \beta(p - p_0) = \alpha_a + \beta(a + c - (a_0 + c_0))$$

$$= \underbrace{\alpha_a + \beta(a - a_0)}_{\text{cohort age-effect}} + \beta(c - c_0)$$

The two **models** are the same.

The **parametrization** is different.

The age-curve refers either

- to a period (cross-sectional rates) or
- to a cohort (longitudinal rates).

Age-drift model (Ad) 151 / 333

Fitting the model in R II


```
> m.apc <- glm( D ~ factor(A) + factor(P) + factor(C) + offset(log(Y)),
+             family = poisson, data = tc )
> summary( m.apc )

Call:
glm(formula = D ~ factor(A) + factor(P) + factor(C) + offset(log(Y)),
    family = poisson, data = tc)

Deviance Residuals:
    Min       1Q   Median       3Q      Max
-1.5406 -0.5534  0.0000  0.4934  1.2966

Coefficients: (1 not defined because of singularities)
            Estimate Std. Error z value Pr(>|z|)
(Intercept) -11.39890   0.23316  -48.889 < 2e-16
factor(A) 22.5    1.19668   0.07789  15.364 < 2e-16
factor(A) 27.5    1.63551   0.08627  18.957 < 2e-16
factor(A) 32.5    1.71939   0.10223  16.819 < 2e-16
factor(A) 37.5    1.57062   0.12205  12.869 < 2e-16
```

Age-Period-Cohort model (APC-cat) 155 / 333



Which age-curve is period and which is cohort?

Age-drift model (Ad) 152 / 333

Fitting the model in R III

```
factor(A) 42.5    1.29418   0.14416  8.977 < 2e-16
factor(A) 47.5    0.87209   0.16828  5.182 2.19e-07
factor(A) 52.5    0.51257   0.19309  2.655 0.00794
factor(A) 57.5    0.12801   0.21109  0.606 0.54424
factor(P) 1950.5  0.20286   0.08247  2.460 0.01390
factor(P) 1955.5  0.42044   0.09081  4.630 3.66e-06
factor(P) 1960.5  0.64099   0.10548  6.077 1.23e-09
factor(P) 1965.5  0.82135   0.12407  6.620 3.60e-11
factor(P) 1970.5  1.06435   0.14444  7.369 1.72e-13
factor(P) 1975.5  1.27796   0.16653  7.674 1.67e-14
factor(P) 1980.5  1.43441   0.18961  7.565 3.88e-14
factor(P) 1985.5  1.50578   0.21339  7.056 1.71e-12
factor(P) 1990.5  1.58798   0.23562  6.740 1.59e-11
factor(C) 1893   0.50556   0.42894  1.179 0.23855
factor(C) 1898   0.56443   0.38398  1.470 0.14158
factor(C) 1903   0.28430   0.35577  0.800 0.42397
factor(C) 1908   0.20683   0.32836  0.630 0.52877
factor(C) 1913   0.22302   0.30344  0.735 0.46236
factor(C) 1918   0.02713   0.28150  0.096 0.92322
```

Age-Period-Cohort model (APC-cat) 156 / 333

Age-Period-Cohort model

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:

Age-Period-Cohort models

— and some cousins

European Doctoral School of Demography,

Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

APC-cat

Fitting the model in R IV

```
factor(C) 1923   0.03280   0.25971  0.126 0.89950
factor(C) 1928   0.02155   0.23945  0.090 0.92830
factor(C) 1933   0.02518   0.21988  0.115 0.90881
factor(C) 1938  -0.07240   0.20268  -0.357 0.72094
factor(C) 1943  -0.35284   0.18706  -1.886 0.05927
factor(C) 1948  -0.30472   0.17308  -1.761 0.07831
factor(C) 1953  -0.17916   0.16258  -1.102 0.27047
factor(C) 1958  -0.11739   0.15585  -0.753 0.45133
factor(C) 1963  -0.10882   0.15410  -0.706 0.48008
factor(C) 1968  -0.16807   0.16235  -1.035 0.30053
factor(C) 1973      NA          NA      NA      NA
```

(Dispersion parameter for poisson family taken to be 1)

Null deviance: 2761.230 on 89 degrees of freedom
Residual deviance: 38.783 on 56 degrees of freedom
AIC: 637.64

Number of Fisher Scoring iterations: 4

Age-Period-Cohort model (APC-cat) 157 / 333

The age-period-cohort model

$$\log[\lambda(a, p)] = \alpha_a + \beta_p + \gamma_c$$

► Three effects:

- a — Age (at diagnosis)
- p — Period (of diagnosis)
- c — Cohort (of birth)

► No assumptions about the **shape** of effects.

► Levels of A , P and C are assumed **exchangeable**

► *i.e.* no assumptions about the relationship between parameter estimates and the **scaled values** of A , P and C

Age-Period-Cohort model (APC-cat) 153 / 333

Fitting the model in R V

Age-Period-Cohort model (APC-cat) 158 / 333

No. of parameters

- A has $9(A)$ levels
- P has $10(P)$ levels
- C=P-A has $18(C = A + P - 1)$ levels
- Age-drift model has $A + 1 = 10$ parameters
- Age-period model has $A + P - 1 = 18$ parameters
- Age-cohort model has $A + C - 1 = 26$ parameters
- Age-period-cohort model has $A + P + C - 3 = 34$ parameters:

```
> length( coef(m.apc) ) ; sum( !is.na(coef(m.apc)) )
[1] 35
[1] 34
```

The missing parameter is because of the **identifiability problem**.


How it all adds up:

$$\begin{aligned} \lambda(a, p) &= \hat{\alpha}_a + \hat{\beta}_p + \hat{\gamma}_c \\ &= \hat{\alpha}_a + \gamma_{c_0} + \hat{\mu}_p + \hat{\delta}_p(a + c_0) + \\ &\quad \hat{\beta}_p - \hat{\mu}_p - \hat{\delta}_p(a + c) + \\ &\quad \hat{\gamma}_c - \gamma_{c_0} + \hat{\delta}_p(c - c_0) \end{aligned}$$

Only the regression on period is needed! (For this model...)

Test for effects

```
> tc.apc <- apc.fit( tc, model="factor", ref.c=1943, scale=10^5 )
[1] "ML of APC-model Poisson with log(Y) offset : ( ACP ): \n"
      Model Mod. df. Mod. dev. Test df. Test dev. Pr(>Chi) Test dev
1      Age      81 1114.65039      NA      NA      NA
2 Age-drift     80 131.77144      1 982.878942 9.457990e-216 982.878
3 Age-Cohort    64  70.20129      16 61.570158 2.840192e-07 3.844
4 Age-Period-Cohort 56  38.78290      8 31.418390 1.183381e-04 3.927
5 Age-Period    72 122.23379      16 83.450895 3.950394e-11 5.215
6 Age-drift     80 131.77144      8  9.537653 2.989863e-01 1.192
```



```
> plot( tc.apc )
```

```
CD.offset RR.fac
Age-Period-Cohort model (APC-cat) 1
```

How to choose a parametrization

- Standard approach: Put extremes of periods or cohorts to 0, and choose a reference for the other.
- Clayton & Schifflers: only 2nd order differences are invariants:

$$\alpha_{i-1} - 2\alpha_i + \alpha_{i+1}$$

Implemented in Epi via the contrast type `contr.2nd` (later).

- Holford: Extract linear effects by regression:

$$\begin{aligned} \lambda(a, p) &= \hat{\alpha}_a + \hat{\beta}_p + \hat{\gamma}_c \\ &= \tilde{\alpha}_a + \hat{\mu}_p + \hat{\delta}_p a + \\ &\quad \tilde{\beta}_p + \hat{\mu}_p + \hat{\delta}_p p + \\ &\quad \tilde{\gamma}_c + \hat{\mu}_c + \hat{\delta}_c c \end{aligned}$$


Customize the frame for nicer plot of parameter estimates:

```
> par( mar=c(3,4,0.1,4), mgp=c(3,1,0)/1.6, las=1 )
> apc.frame( a.lab=c(2,4,6)*10,
+           a.tic=1:6*10,
+           cp.lab=1900+0:4*20,
+           cp.tic=1890+0:10*10,
+           r.lab=c(1,2,5), c(1,2)*10,
+           r.tic=c(1:10,15,20,25),
+           rr.ref=5 )
> matshade( tc.apc$Age[,1], tc.apc$Age[,-1], lwd=2 )
> pc.matshade( tc.apc$Per[,1], tc.apc$Per[,-1], lwd=2 )
> pc.matshade( tc.apc$Coh[,1], tc.apc$Coh[,-1], lwd=2 )
> pc.points( 1943, 1, pch=16 )
```

Assumptions

Assumptions are needed to do this, e.g.:

- Age is the major time scale
- Cohort is the secondary time scale (the major secular trend)
- c_0 is the reference cohort
- Period is the residual time scale: 0 on average, 0 slope
- ... constraining first and last period parameter to 0 is a crude way of obtaining this.



Relocating effects between A, P and C

Period effect, 0 on average, slope is 0: a regression of β_p on p :

$$g(p) = \tilde{\beta}_p = \beta_p - \hat{\mu}_p - \hat{\delta}_p p$$

Cohort effect, absorbing all time-trend ($\delta_p p = \delta_p(a + c)$) and risk relative to c_0 :

$$h(c) = \gamma_c - \gamma_{c_0} + \hat{\delta}_p(c - c_0)$$

The rest is the age-effect:

$$f(a) = \alpha_a + \hat{\mu}_p + \hat{\delta}_p a + \hat{\delta}_p c_0 + \gamma_{c_0}$$

A simple practical approach

- First fit the age-cohort model, with cohort c_0 as reference and get estimates $\hat{\alpha}_a$ and $\hat{\gamma}_c$:

$$\log[\lambda(a, p)] = \hat{\alpha}_a + \hat{\gamma}_c$$

- Then consider the full APC-model with age and cohort effects constrained to be as estimated from the AC-model:

$$\log[\lambda(a, p)] = \hat{\alpha}_a + \hat{\gamma}_c + \beta_p$$

- ▶ The residual period effect can be estimated if we note that for the number of cases we have:

$$\log(\text{expected cases}) = \log[\hat{\lambda}(a, p) Y] = \underbrace{\hat{\alpha}_a + \hat{\gamma}_c + \log(Y)}_{\text{"known"}} + \beta_p$$

- ▶ This is analogous to the expression for a Poisson model in general,
- ▶ ... but now is the offset not just $\log(Y)$ but $\hat{\alpha}_a + \hat{\gamma}_c + \log(Y)$, the log of the fitted values from the age-cohort model.
- ▶ β_p s are estimated in a Poisson model with this as offset.
- ▶ Advantage: We get the standard errors for free.

Age at entry (diagnosis) as covariate

t : time since entry (duration)

e : age at entry

$a = e + t$: current age (age at follow-up)

Duration as basic time-scale; linear effect of age at entry:

$$\log(\lambda(a, t)) = f(t) + \beta e = (f(t) - \beta t) + \beta a$$

Immaterial whether a or e is used as (log)-**linear** covariate as long as t is in the model.

Customize the frame for nicer plot of parameter estimates:

```
> par( mar=c(3,4,0.1,4), mgp=c(3,1,0)/1.6, las=1 )
> apc.frame( a.lab=c(2,4,6)*10,
+           a.tic=1:6*10,
+           cp.lab=1900+0:4*20,
+           cp.tic=1890+0:10*10,
+           r.lab=c(c(1,2,5),c(1,2)*10),
+           r.tic=c(1:10,15,20,25),
+           rr.ref=5 )
+ matshade( tc.acp$Age[,1], tc.acp$Age[,-1], lwd=2, alpha=0.2 )
> pc.matshade( tc.acp$Per[,1], tc.acp$Per[,-1], lwd=2, alpha=0.2 )
> pc.matshade( tc.acp$Coh[,1], tc.acp$Coh[,-1], lwd=2, alpha=0.2 )
> pc.points( 1943, 1, pch=16 )
> #
> # The stepwise conditioning:
> tc.ac.p <- apc.fit( tc, model="factor", parm="AC-P", ref.c=1943, scale=10^5 )
```

Non-linear effects of time-scales

Arbitrary effects of the three variables t , a and e :
 \Rightarrow genuine extension of the model.

$$\log(\lambda(a, t)) = f(t) + g(a) + h(e)$$

Three quantities can be arbitrarily moved between the three functions:

$$\tilde{f}(t) = f(t) - \mu_a - \mu_e + \gamma t$$

$$\tilde{g}(a) = g(a) + \mu_a - \gamma a$$

$$\tilde{h}(e) = h(e) + \mu_e + \gamma e$$

because $t - a + e = 0$.

This is the age-period-cohort modeling problem again.


```
[1] "Sequential modelling Poisson with log(Y) offset : ( AC-P ):\n"
      Model Mod. df. Mod. dev. Test df. Test dev. Pr(>Chi) Test dev.
1      Age      81 1114.65039      NA      NA      NA      NA
2 Age-drift    80 131.77144      1 982.878942 9.457990e-216 982.878
3 Age-Cohort   64 70.20129      16 61.570158 2.840192e-07 3.848
4 Age-Period  56 38.78290      8 31.418390 1.183381e-04 3.927
5 Age-Period  72 122.23379      16 83.450895 3.950394e-11 5.214
6 Age-drift    80 131.77144      8 9.537653 2.989863e-01 1.192

> #
> matshade( tc.ac.p$Age[,1], tc.ac.p$Age[,-1], lwd=2, alpha=0.2, lty='22', len=
> pc.matshade( tc.ac.p$Per[,1], tc.ac.p$Per[,-1], lwd=2, alpha=0.2, lty='22', len=
> pc.matshade( tc.ac.p$Coh[,1], tc.ac.p$Coh[,-1], lwd=2, alpha=0.2, lty='22', len=
```

"Controlling for age"

— is not a well defined statement:

- ▶ Mostly it means that age **at entry** is included in the model.
- ▶ But ideally one would check whether there were non-linear effects of age at entry and current age.
- ▶ This would require modeling of multiple timescales.
- ▶ Which is best accomplished by splitting follow up and using Poisson models, with time scales as covariates.



Tabulation in the Lexis diagram

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:

Age-Period-Cohort models

— and some cousins

European Doctoral School of Demography,

Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

Age at entry Age-Duration-Diagnosis

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:

Age-Period-Cohort models

— and some cousins

European Doctoral School of Demography,

Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>


Tabulation of register data

Age	1943	1953	1963	1973	1983	1993										
65	471.0	512.8	571.1	622.5	680.8	696.2	728	683.8	43	686.4	42	940.9	34	627.7	45	544.8
55	539.4	600.3	653.9	715.4	732.7	718.3	724.2	675.5	660.8	64	721.1	64	51	701.5		
45	622.1	676.7	737.9	753.5	738.1	746.4	689.2	682.4	743.1	52	86	96	96	817.8		
35	65	41	65	64	67	85	103	119	121	121	155	126				
25	694.1	754.3	785.5	749.9	765.5	709.8	695.5	727.8	945.3	1035.7	754.5					
15	759.3	774.5	769.3	711.6	700.1	769.9	965.4	1045.3	955.0	967.1	821.5					
5	790.5	781.8	723.0	688.6	784.8	962.7	1056.1	960.9	966.2	1031.6	835.7					
0	813.0	744.7	721.8	770.9	960.3	1053.8	967.5	953.0	1019.7	1017.3	760.9					
-5	10	7	13	13	15	33	35	37	49	51	41					
-15	773.8	744.2	794.1	972.9	1051.5	961.0	952.5	1011.1	1005.0	929.8	670.2					

Testis cancer cases in Denmark.

Male person-years in Denmark.

Tabulation by age, period and cohort




Gives triangular sets with differing mean age, period and cohort:

These correct midpoints for age, period and cohort must be used in modeling.

A person dying in age a at date p in **B** contributes a risk time in **B**, so the average will be:

$$\int_{p=0}^{p=1} \int_{a=0}^{a=p} 2a \, da \, dp = \int_{p=0}^{p=1} p^2 \, dp = \frac{1}{3}$$




From population figures to risk time

Population figures in the form of size of the population at certain date are available from most statistical bureaus.

This corresponds to population sizes along the vertical lines in the diagram.

We want risk time figures for the population in the squares and triangles in the diagram.



Mean contributions to risk time in A and B:


	A:	B:
Survivors:	$\ell_{a+1,p+1} \times \frac{1}{2}y$	$\ell_{a+1,p+1} \times \frac{1}{2}y$
Dead in A :	$\frac{1}{2}(\ell_{a,p} - \ell_{a+1,p+1}) \times \frac{1}{3}y$	
Dead in B :	$\frac{1}{2}(\ell_{a,p} - \ell_{a+1,p+1}) \times \frac{1}{3}y$	$\frac{1}{2}(\ell_{a,p} - \ell_{a+1,p+1}) \times \frac{1}{3}y$
Σ	$(\frac{1}{3}\ell_{a,p} + \frac{1}{6}\ell_{a+1,p+1}) \times y$	$(\frac{1}{6}\ell_{a,p} + \frac{1}{3}\ell_{a+1,p+1}) \times y$

The number of deaths in **A** and **B** is $\ell_{a,p} - \ell_{a+1,p+1}$, and we assume that half occur in **A** and half in **B**.

Prevalent population figures

$\ell_{a,p}$ is the number of persons in age class a alive at the beginning of period (=year) p .

The aim is to compute person-years for the triangles **A** and **B**, respectively.




Population as of 1. January from Statistics Denmark:

Age	Men			Women		
	2000	2001	2002	2000	2001	2002
22	33435	33540	32272	32637	32802	31709
23	35357	33579	33742	34163	32853	33156
24	38199	35400	33674	37803	34353	33070
25	37958	38257	35499	37318	37955	34526
26	38194	38048	38341	37292	37371	38119
27	39891	38221	38082	39273	37403	37525

The area of the triangle is $1/2$, so the uniform measure over the triangle has density 2. Therefore a person dying in age a at date p in **A** contributes p risk time in **A**, so the average will be:


$$\int_{p=0}^{p=1} \int_{a=p}^{a=1} 2p \, da \, dp = \int_{p=0}^{p=1} 2p - 2p^2 \, dp = \left[p^2 - \frac{2p^3}{3} \right]_{p=0}^{p=1} = \frac{1}{3}$$



Exercise:


Fill in the risk time figures in as many triangles as possible from the previous table for men and women, respectively.

Look at the **N2Y** function in **Epi**.



A person dying in age a at date p in **B** contributes $p - a$ risk time in **A**, so the average will be (again using the density 2 of the uniform measure):

$$\int_{p=0}^{p=1} \int_{a=0}^{a=p} 2(p-a) \, da \, dp = \int_{p=0}^{p=1} [2pa - a^2]_{a=0}^{a=p} \, dp = \int_{p=0}^{p=1} p^2 \, dp = \frac{1}{3}$$




Summary:

Population risk time:

$$\mathbf{A:} \left(\frac{1}{3}\ell_{a,p} + \frac{1}{6}\ell_{a+1,p+1} \right) \times y$$

$$\mathbf{B:} \left(\frac{1}{6}\ell_{a-1,p} + \frac{1}{3}\ell_{a,p+1} \right) \times y$$

Mean age, period and cohort: $\frac{1}{3}$ into the interval.



APC-model for triangular data

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020


<http://BendixCarstensen.com/APC/EDSD-2020>

APC-tri

Fill in the number of cases (D) and person-years (Y) from previous slide.

Indicate birth cohorts on the axes for upper and lower triangles.

Mark mean date of birth for these.



APC-model for triangular data (APC-tri)

201/ 333

Model for triangular data

- ▶ One parameter per distinct value on each timescale.
- ▶ Example: 3 age-classes and 3 periods:
 - ▶ 6 age parameters
 - ▶ 6 period parameters
 - ▶ 10 cohort parameters
- ▶ Model:

$$\lambda_{ap} = \alpha_a + \beta_p + \gamma_c$$

APC-model for triangular data (APC-tri)

197/ 333

APC-model with “synthetic” cohorts

```
> mc <- glm( D ~ factor(A5) - 1 +
+           factor(P5-A5) +
+           factor(P5) + offset( log( Y ) ),
+           family=poisson )
> summary( mc )

...

Null deviance: 1.0037e+08 on 220 degrees of freedom
Residual deviance: 8.8866e+02 on 182 degrees of freedom
```

No. parameters: 220 – 182 = 38.

$$A = 10, \quad P = 11, \quad C = 20 \quad \Rightarrow \quad A + P + C - 3 = 38$$

APC-model for triangular data (APC-tri)

202/ 333

Problem: Disconnected design!

Log-likelihood contribution from one triangle:

$$D_{ap} \log(\lambda_{ap}) - \lambda_{ap} Y_{ap} = D_{ap} \log(\alpha_a + \beta_p + \gamma_c) - (\alpha_a + \beta_p + \gamma_c) Y_{ap}$$

The log-likelihood can be separated:

$$\sum_{a,p \in \nabla} D_{ap} \log(\lambda_{ap}) - \lambda_{ap} Y_{ap} + \sum_{a,p \in \triangle} D_{ap} \log(\lambda_{ap}) - \lambda_{ap} Y_{ap}$$

No common parameters between terms
— we have two separate models:
One for upper triangles, one for lower.

APC-model for triangular data (APC-tri)

198/ 333

APC-model with “correct” cohorts

```
> mx <- glm( D ~ factor(Ax) - 1 +
+           factor(Cx) +
+           factor(Px) + offset( log( Y ) ),
+           family=poisson )
> summary( mx )

...

Null deviance: 1.0037e+08 on 220 degrees of freedom
Residual deviance: 2.8473e+02 on 144 degrees of freedom
```

No. parameters: 220 – 144 = 76 (= 38 × 2).

$$A = 20, \quad P = 22, \quad C = 40 \quad \Rightarrow \quad A + P + C - 3 = 79 \neq 76!$$

We have fitted two age-period-cohort models separately to upper and lower triangles.

APC-model for triangular data (APC-tri)

203/ 333

Illustration by lung cancer data

```
> library( Epi )
> data( lungDK )
> lungDK[1:10,]
  A5  P5  C5 up   Ax      Px      Cx  D      Y
1 40 1943 1898 1 43.33333 1944.667 1901.333 52 336233.8
2 40 1943 1903 0 41.66667 1946.333 1904.667 28 357812.7
3 40 1948 1903 1 43.33333 1949.667 1906.333 51 363783.7
4 40 1948 1908 0 41.66667 1951.333 1909.667 30 390985.8
5 40 1953 1908 1 43.33333 1954.667 1911.333 50 391925.3
6 40 1953 1913 0 41.66667 1956.333 1914.667 23 377515.3
7 40 1958 1913 1 43.33333 1959.667 1916.333 56 365575.5
8 40 1958 1918 0 41.66667 1961.333 1919.667 43 383689.0
9 40 1963 1918 1 43.33333 1964.667 1921.333 44 385878.5
10 40 1963 1923 0 41.66667 1966.333 1924.667 38 371361.5
```


APC-model for triangular data (APC-tri)

199/ 333

Fill in the number of cases (D) and person-years (Y) from previous slide.


Indicate birth cohorts on the axes for upper and lower triangles.

Mark mean date of birth for these.




APC-model for triangular data (APC-tri)

200/ 333



APC-model for triangular data (APC-tri)

204/ 333



APC-model for triangular data (APC-tri)

205/ 333

Now, explicitly fit models for upper and lower triangles:


```
> mx.u <- glm( D ~ factor(Ax) - 1 +
+             factor(Cx) +
+             factor(Px) + offset( log( Y/10^5 ) ), family=poisson,
+             data=lungDK [lungDK$up=1,] )
> mx.l <- glm( D ~ factor(Ax) - 1 +
+             factor(Cx) +
+             factor(Px) + offset( log( Y/10^5 ) ), family=poisson,
+             data=lungDK [lungDK$up=0,] )
> mx.u$deviance
[1] 284.7269
> mx.l$deviance
[1] 134.4566
> mx.u$deviance
[1] 150.2703
> mx.l$deviance+mx.u$deviance
[1] 284.7269
```

Non-linear effects

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>



Testis cancer

Testis cancer in Denmark:

```
> library( Epi )
> data( testisDK )
> str( testisDK )

'data.frame': 4860 obs. of 4 variables:
 $ A: num 0 1 2 3 4 5 6 7 8 9 ...
 $ P: num 1943 1943 1943 1943 1943 ...
 $ D: num 1 1 0 1 0 0 0 0 0 0 ...
 $ Y: num 39650 36943 34588 33267 32614 ...

> head( testisDK )
  A  P  D  Y
1 0 1943 1 39649.50
2 1 1943 1 36942.83
3 2 1943 0 34588.33
4 3 1943 1 33267.00
5 4 1943 0 32614.00
6 5 1943 0 32020.33
```

Modeling for Lexis triangles

- ▶ Modeling by factors not possible
- ▶ Two separate models that cannot be fitted together
- ▶ We are not using the **quantitative** values of age, period and cohort.
- ▶ **Solution:** parametric models using the quantitative nature of a , p and $c = p - a$.
- ▶ ... so we need to handle smooth parametric functions.

Cases, PY and rates

```
> print(
+ stat.table( list( A = floor(A/10)*10,
+                 P = floor(P/10)*10,
+                 list( D = sum(D),
+                       Y = sum(Y/1000),
+                       rate = ratio(D,Y,10^6) ),
+                 margins = TRUE, data = testisDK ), digits=c(sum=0, ratio=2) )
```

A	P						Total
	1940	1950	1960	1970	1980	1990	
0	10 2605 3.84	7 4037 1.73	16 3885 4.12	18 3821 4.71	9 3071 2.93	10 2166 4.62	70 19584 3.57
10	13 2136 6.09	27 3505 7.70	37 4004 9.24	72 3906 18.43	97 3847 25.21	75 2261 33.17	321 19659 16.33
20	124 2026	221 2622	280 2402	535 4020	724 2944	557 2825	2441 10245

Tuesday exercise: Age-drift model

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

Linear effects in glm

How do rates depend on age?

```
> ml <- glm( D ~ A, offset=log(Y), family=poisson, data=testisDK )
> round( ci.lin( ml ), 4 )

              Estimate StdErr      z P      2.5% 97.5%
(Intercept) -9.7755 0.0207 -472.3164 0 -9.8160 -9.7349
A             0.0055 0.0005  11.3926 0  0.0045  0.0064

> round( ci.exp( ml ), 4 )

              exp(Est.) 2.5% 97.5%
(Intercept)  0.0001 0.0001 0.0001
A             1.0055 1.0046 1.0064
```

Linear increase of log-rates by age

Tuesday afternoon exercise

1. Load the `Epi` package.
2. Do the practical 2.4 (Age-drift model). These are both prerequisites for APC-models.
3. Most of Wednesday will be concerned with exercise 2.5
4. I will be available the first approx. hour for direct questions that will be broadcast to the entire audience.
5. After that I will be on e-mail, until about 15.
6. Tuesday morning we will:
 - ▶ Recap the contents of today's lecture
 - ▶ Go over the solutions to today's exercises
7. Remember: Questions always welcome at any time, just switch on you microphone and peak.


Linear effects in glm

```
> nd <- data.frame( A=15:60, Y=10^5 )
> pr <- ci.pred( ml, newdata=nd )
> head( pr )

              Estimate 2.5% 97.5%
1 6.170105 5.991630 6.353896
2 6.204034 6.028525 6.384652
3 6.238149 6.065547 6.415662
4 6.272452 6.102689 6.446937
5 6.306943 6.139944 6.478485
6 6.341624 6.177301 6.510319

> matplot( nd$A, pr, type="l", lty=1, lwd=c(3,1), col="black", log="y" )
```

Linear effects in glm



```
> nd <- data.frame( A=15:60, Y=10^5 )
> pr <- ci.pred( ml, newdata=nd )
> matshade( nd$A, pr, plot=TRUE, lwd=3, log="y" )
```

Non-linear effects (crr-mod)

214/ 333

Adding a linear period effect

```
> msp <- glm( D ~ Ns(A,knots=seq(15,65,10)) + P,
+ offset=log(Y), family=poisson, data=testisDK )
> nd <- data.frame( A=15:60, Y=10^5, P=1970 )
```

A multiplicative model:

$$\lambda(a, p) = f(a) \times g(p), \quad g(p_{ref}) = 1$$

$f(a)$: Rate at p_{ref}

$g(p)$: Rate ratio relative to p_{ref}

Non-linear effects (crr-mod)

219/ 333

Quadratic effects in glm

How do rates depend on age?

```
> mq <- glm( D ~ A + I(A^2),
+ offset=log(Y), family=poisson, data=testisDK )
> round( ci.lin( mq ), 4 )
```

	Estimate	StdErr	z	P	2.5%	97.5%
(Intercept)	-12.3656	0.0596	-207.3611	0	-12.4825	-12.2487
A	0.1806	0.0033	54.8290	0	0.1741	0.1871
I(A^2)	-0.0023	0.0000	-53.7006	0	-0.0024	-0.0022


```
> round( ci.exp( mq ), 4 )
```

	exp(Est.)	2.5%	97.5%
(Intercept)	0.0000	0.0000	0.0000
A	1.1979	1.1902	1.2057
I(A^2)	0.9977	0.9976	0.9978

Non-linear effects (crr-mod)

215/ 333

Adding a linear period effect




```
> matshade( nd$A, ci.pred(msp,nd), plot=TRUE,
+ log="y", xlab="Age", ylim=c(2,20), col="brown", alpha=0.15,
+ ylab="Testis cancer incidence rate per 100,000 PY in 1970" )
```

Non-linear effects (crr-mod)

220/ 333

Quadratic effect in glm



```
> matshade( nd$A, cbind( ci.pred(mq,nd), ci.pred(ml,nd) ), plot=TRUE,
+ log="y", col=c("blue","black"), alpha=c(15,5)/100 )
```

Non-linear effects (crr-mod)

216/ 333

The period effect

```
> nd.p <- data.frame( P=1945:1995 )
> nd.r <- data.frame( P=1970 )
> str( nd.p )
```

```
'data.frame': 51 obs. of 1 variable:
 $ P: int 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 ...
```

```
> str( nd.r )
```

```
'data.frame': 1 obs. of 1 variable:
 $ P: num 1970
```

```
> RR <- ci.exp( msp, ctr.mat=list(nd.p,nd.r), xvars="A" )
> matshade( nd.p$P, RR, plot=TRUE,
+ log="y", xlab="Date", ylab="Testis cancer incidence RR",
+ type="l", lty=1, lwd=c(3,1,1), col="black" )
> abline( v=1970, h=1, col="red" )
```

Non-linear effects (crr-mod)

221/ 333

Spline effects in glm

```
> library( splines )
> ms <- glm( D ~ Ns(A,knots=seq(15,65,10)),
+ offset=log(Y), family=poisson, data=testisDK )
> round( ci.exp( ms ), 3 )
```

	exp(Est.)	2.5%	97.5%
(Intercept)	0.000	0.000	0.000
Ns(A, knots = seq(15, 65, 10))1	8.548	7.650	9.551
Ns(A, knots = seq(15, 65, 10))2	5.706	4.998	6.514
Ns(A, knots = seq(15, 65, 10))3	1.002	0.890	1.128
Ns(A, knots = seq(15, 65, 10))4	14.402	11.896	17.436
Ns(A, knots = seq(15, 65, 10))5	0.466	0.429	0.505

```
> matplot( nd$A, ci.pred( ms, nd ),
+ log="y", xlab="Age", ylab="Testis cancer incidence rate per 100,000 PY",
+ type="l", lty=1, lwd=c(3,1,1), col="black", ylim=c(2,20) )
```

Non-linear effects (crr-mod)

217/ 333

A quadratic period effect


```
> mspq <- glm( D ~ Ns(A,knots=seq(15,65,10)) + P + I(P^2),
+ offset=log(Y), family=poisson, data=testisDK )
> round( ci.exp( mspq ), 4 )
```

	exp(Est.)	2.5%	97.5%
(Intercept)	0.0000	0.0000	0.0000
Ns(A, knots = seq(15, 65, 10))1	8.3560	7.4783	9.3366
Ns(A, knots = seq(15, 65, 10))2	5.5133	4.8290	6.2945
Ns(A, knots = seq(15, 65, 10))3	1.0060	0.8935	1.1326
Ns(A, knots = seq(15, 65, 10))4	13.4388	11.1008	16.2691
Ns(A, knots = seq(15, 65, 10))5	0.4582	0.4223	0.4971
P	2.1893	1.4566	3.2906
I(P^2)	0.9998	0.9997	0.9999

Non-linear effects (crr-mod)

222/ 333

Spline effects in glm




```
> matshade( nd$A, cbind( ci.pred(ms,nd), ci.pred(mq,nd) ), plot=TRUE,
+ log="y", xlab="Age", ylab="Testis cancer incidence rate per 100,000 PY",
+ col=c("brown","blue"), alpha=c(15,10)/100, ylim=c(2,20) )
```

Non-linear effects (crr-mod)

218/ 333

A quadratic period effect



```
> matshade( nd.p$P, ci.exp( mspq, ctr.mat=list(nd.p,nd.r), xvars="A" ), plot=TRUE,
+ log="y", xlab="Date", ylab="Testis cancer incidence RR", col="blue" )
> abline( h=1, v=1970, col="red", lty="13" )
```

Non-linear effects (crr-mod)

223/ 333

A spline period effect

```
> msp <- glm( D ~ Ns(A,knots=seq(15,65,10)) +
+           Ns(P,knots=seq(1950,1990,10)),
+           offset=log(Y), family=poisson, data=testisDK )
> round( ci.exp( msp ), 3 )
```

	exp(Est.)	2.5%	97.5%
(Intercept)	0.000	0.000	0.000
Ns(A, knots = seq(15, 65, 10))1	8.327	7.452	9.305
Ns(A, knots = seq(15, 65, 10))2	5.528	4.842	6.312
Ns(A, knots = seq(15, 65, 10))3	1.007	0.894	1.133
Ns(A, knots = seq(15, 65, 10))4	13.447	11.107	16.279
Ns(A, knots = seq(15, 65, 10))5	0.458	0.422	0.497
Ns(P, knots = seq(1950, 1990, 10))1	1.711	1.526	1.918
Ns(P, knots = seq(1950, 1990, 10))2	2.190	2.028	2.364
Ns(P, knots = seq(1950, 1990, 10))3	3.222	2.835	3.661
Ns(P, knots = seq(1950, 1990, 10))4	2.299	2.149	2.459


Wednesday exercise 1: Linear and curved effects

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

Period effect



```
> matshade( nd.p$P, ci.exp( msp, ctr.mat=list(nd.p,nd.r), xvars="A" ), plot=TRUE
+           log="y", xlab="Date", ylab="Testis cancer incidence RR", lwd=3 )
> abline( h=1, v=1970 )
```

Wednesday exercise

1. Load the `Epi` package.
2. Do the practical 2.3: Linear and curved effects.
3. In about 20 min min I will be back and we shall go over the exercise.

Period effect

```
> par( mfrow=c(1,2) )
> matshade( nd$A, ci.pred( msp, nd ), plot=TRUE,
+           log="y", xlab="Age", col="black",
+           ylab="Testis cancer incidence rate per 100,000 PY in 1970" )
> matshade( nd.p$P, ci.exp( msp, ctr.mat=list(nd.p,nd.r), xvars="A" ), plot=TRUE
+           log="y", xlab="Date", ylab="Testis cancer incidence RR",
+           col="black" )
> abline( h=1, v=1970 )
```


APC-model: Parametrization

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

Age and period effect



What's the problem?

- ▶ One parameter is assigned to each distinct value of the timescales, the **scale** of the variables is not used.
- ▶ The solution is to "tie together" the points on the scales together with smooth functions of the **mean** age, period and cohort with three functions:

$$\lambda_{ap} = f(a) + g(p) + h(c)$$

- ▶ The practical problem is how to choose a reasonable parametrization of these functions, and how to get estimates.

Age and period effect with `ci.exp`

- ▶ In rate models there is always one term with the **rate** dimension.
Usually **age**
- ▶ But it must refer to specific **reference** values for **all other** variables (in this case only P).
- ▶ For the "other" variables, report the RR **relative** to the reference point.
- ▶ Only parameters relevant for the variable (P) actually used in the calculation.
- ▶ We are computing the difference between two predictions.
- ▶ ... as well as the confidence intervals for it.

The identifiability problem still exists:

$$c = p - a \Leftrightarrow p - a - c = 0$$

$$\begin{aligned} \lambda_{ap} &= f(a) + g(p) + h(c) \\ &= f(a) + g(p) + h(c) + \gamma(p - a - c) \\ &= f(a) - \mu_a - \gamma a + \\ &\quad g(p) + \mu_a + \mu_c + \gamma p + \\ &\quad h(c) - \mu_c - \gamma c \end{aligned}$$

A decision on parametrization is needed.
... it must be **external to the model**.

Smooth functions

$$\log(\lambda(a, p)) = f(a) + g(p) + h(c)$$

Possible choices for non-linear parametric functions describing the effect of the three **quantitative** variables:

- ▶ Polynomials / fractional polynomials.
- ▶ Linear / quadratic / cubic splines.
- ▶ Natural splines.

All of these contain the linear effect as special case.

“Extract the trend”

- ▶ **Not** a well-defined concept:
 - ▶ Regress $\hat{g}(p)$ on p for all units in the dataset.
 - ▶ Regress $\hat{g}(p)$ on p for all different values of p .
 - ▶ Weighted regression — what weights?
- ▶ How do we get the standard errors?
- ▶ Matrix-algebra!
- ▶ Projections!
- ▶ Weighted inner product...

Parametrization of effects

There are still three “free” parameters:

$$\tilde{f}(a) = f(a) - \mu_a - \gamma a$$

$$\tilde{g}(p) = g(p) + \mu_a + \mu_c + \gamma p$$

$$\tilde{h}(c) = h(c) - \mu_c - \gamma c$$

Any set of 3 numbers, μ_a , μ_c and γ will produce effects with the same sum:

$$\tilde{f}(a) + \tilde{g}(p) + \tilde{h}(c) = f(a) + g(p) + h(c)$$

The problem is to choose μ_a , μ_c and γ according to some criterion for the functions.

Parametric function

Suppose that $g(p)$ is parametrized using the design matrix \mathbf{M} , with the estimated parameters π .

Example: 2nd degree polynomial:

$$\mathbf{M} = \begin{bmatrix} 1 & p_1 & p_1^2 \\ 1 & p_2 & p_2^2 \\ \vdots & \vdots & \vdots \\ 1 & p_n & p_n^2 \end{bmatrix} \quad \pi = \begin{bmatrix} \pi_0 \\ \pi_1 \\ \pi_2 \end{bmatrix} \quad g(p) = \mathbf{M}\pi$$

$\text{nrow}(\mathbf{M})$ is the no. of observations in the dataset,

$\text{ncol}(\mathbf{M})$ is the no. of parameters

Parametrization principle

1. The age-function should be interpretable as log age-specific rates in a cohort c_0 after adjustment for the period effect.
2. The cohort function is 0 at a reference cohort c_0 , interpretable as log-RR relative to cohort c_0 .
3. The period function is 0 on average with 0 slope, interpretable as log-RR relative to the age-cohort prediction. (residual log-RR).

This will yield cohort age-effects a.k.a. **longitudinal** age effects.

Biologically interpretable:

— what happens during the lifespan of a cohort?

Extract the trend from g :

Vectors \mathbf{x} and \mathbf{y} are orthogonal if the inner product is 0

$$\mathbf{x} \perp \mathbf{y} \Leftrightarrow \langle \mathbf{x} | \mathbf{y} \rangle = \sum_i x_i y_i = 0$$

- ▶ $\langle \tilde{g}(p) | 1 \rangle = 0$, $\langle \tilde{g}(p) | p \rangle = 0$, i.e. \tilde{g} is **orthogonal** to $[1:p]$.
- ▶ Suppose $\tilde{g}(p) = \tilde{\mathbf{M}}\pi$, then for **any** parameter vector π :

$$\langle \tilde{\mathbf{M}}\pi | 1 \rangle = 0, \quad \langle \tilde{\mathbf{M}}\pi | p \rangle = 0 \quad \Rightarrow \quad \tilde{\mathbf{M}} \perp [1:p]$$
- ▶ Thus we just need to be able to produce $\tilde{\mathbf{M}}$ from \mathbf{M} :
Projection on the orthogonal complement of $\text{span}([1:p])$.
- ▶ **But:** orthogonality requires an inner product!

Period-major parametrization

- ▶ Alternatively, the period function could be constrained to be 0 at a reference date, p_0 .
- ▶ Then, age-effects at $a_0 = p_0 - c_0$ would equal the fitted rate for period p_0 (and cohort c_0), and the period effects would be residual log-RRs relative to p_0 .
- ▶ Gives period or **cross-sectional** age-effects
- ▶ Bureaucratically interpretable:
— what was seen at a particular date?

Practical parametrization

1. Set up model matrices for age, period and cohort, M_a , M_p and M_c . Intercept in all three.
2. Extract the linear trend from M_p and M_c , by projecting their columns onto the orthogonal complement of $[1:p]$ and $[1:c]$, respectively
3. Center the cohort effect around c_0 :
Take a row from \tilde{M}_c corresponding to c_0 , replicate to dimension as \tilde{M}_c , and subtract it from \tilde{M}_c to form \tilde{M}_{c_0} .

Implementation:

1. Obtain any set of parameters $f(a)$, $g(p)$, $h(c)$.
2. Extract the trend from the period effect (find μ and β):

$$\tilde{g}(p) = \hat{g}(p) - (\mu + \beta p)$$

3. Decide on a reference cohort c_0 .

4. Use the functions:

$$\tilde{f}(a) = \hat{f}(a) + \mu + \beta a + \hat{h}(c_0) + \beta c_0$$

$$\tilde{g}(p) = \hat{g}(p) - \mu - \beta p$$

$$\tilde{h}(c) = \hat{h}(c) + \beta c - \hat{h}(c_0) - \beta c_0$$

4. Use:
 - M_a for the age-effects,
 - \tilde{M}_p for the period effects and
 - $[c - c_0; \tilde{M}_{c_0}]$ for the cohort effects.
5. Value of $\hat{f}(a)$ is $M_a \hat{\beta}_a$, similarly for the other two effects. Variance is found by $M_a' \hat{\Sigma}_a M_a$, where $\hat{\Sigma}_a$ is the variance-covariance matrix of $\hat{\beta}_a$.

Information about a parameter in the data

Information about log-rate $\theta = \log(\lambda)$:

$$l(\theta|D, Y) = D\theta - e^\theta Y, \quad l'_\theta = D - e^\theta Y, \quad l''_\theta = -e^\theta Y$$

$$\text{so } I(\hat{\theta}) = e^{\hat{\theta}} Y = \hat{\lambda} Y = D.$$

Information about square root of rate $\sigma = \sqrt{\lambda}$:

$$l(\sigma|D, Y) = D\log(\sigma^2) - \sigma^2 Y, \quad l'_\sigma = (D/\sigma^2)2\sigma - 2\sigma Y = 2D/\sigma - 2\sigma Y$$

$$l''_\sigma = -2D/\sigma^2 - 2Y$$

$$\text{so } I(\hat{\sigma}) = -2D/\hat{\sigma}^2 - 2Y = -4Y$$

Consult the help page for: `apc.fit` to see options for weights in inner product, type of function, variants of parametrization etc.

`apc.plot`, `apc.lines` and `apc.frame` to see how to plot the results.


Information in the data and inner product

► Inner products:

$$\langle \mathbf{m}_j | \mathbf{m}_k \rangle = \sum_i m_{ij} m_{ik} \quad \langle \mathbf{m}_j | \mathbf{m}_k \rangle = \sum_i m_{ij} w_i m_{ik}$$

► Weights could be chosen as:

- $w_i = Y_i$, i.e. proportional to the information content for $\sigma = \sqrt{\lambda}$, `dr.extr = Y` (the default)
- $w_i = D_i$, i.e. proportional to the information content for $\theta = \log(\lambda)$, `dr.extr ∈ c(D, T)`
- $w_i = Y_i^2/D_i$, i.e. proportional to the information content for λ , `dr.extr ∈ c(L, R)`
- $w_i = 1$, the "usual" inner product — implicitly used in most of the literature — any other (character) value for `dr.extr`.



How to? I

Implemented in `apc.fit` in the `Epi` package:

```
> library( Epi )
> library( splines )
> data( lungDK )

> mw <- apc.fit( A = lungDK$Ax,
+               P = lungDK$Px,
+               D = lungDK$D,
+               Y = lungDK$Y/1000,
+               ref.c = 1900,
+               npar = 8,
+               parm = "ACP",
+               dr.extr = "y", print.ADV=FALSE ) # drift extraction - choice of inner
```

Other models I

```
> lungDK$A <- lungDK$Ax
> lungDK$P <- lungDK$Px
> ml <- apc.fit( data = lungDK,
+               npar = 8,
+               ref.c = 1900,
+               dr.extr = "y" )

NOTE: npar is specified as:
A P C
8 8 8
[1] "ML of APC-model Poisson with log(Y) offset : ( ACP ):\n"
      Model Mod. df. Mod. dev. Test df. Test dev. Pr(>Chi) Test dev/di
1      Age      212 15468.603      NA      NA      NA      NA
2      Age-drift 211 6858.883      1 8609.7199 0.00000e+00 8609.7199
3      Age-Cohort 205 1034.737      6 5824.1456 0.00000e+00 970.6909
4 Age-Period-Cohort 199 423.158      6 611.5791 7.41219e-129 101.9298
5      Age-Period 205 3082.602      6 2659.4439 0.00000e+00 443.2407
6      Age-drift 211 6858.883      6 3776.2808 0.00000e+00 629.3801
```

How to? II

NOTE: npar is specified as:

```
A P C
8 8 8

> mw$Ref
Per Coh
NA 1900

> cbind( mw$Age[1:4,1:2], mw$Per[1:4,1:2], mw$Coh[1:4,1:2] )
      Age Rate Per P-RR Coh C-RR
[1,] 41.66667 0.08648277 1944.667 0.8584085 1856.333 0.04890071
[2,] 43.33333 0.11507077 1946.333 0.8736385 1859.667 0.06214798
[3,] 46.66667 0.20372101 1949.667 0.9049139 1861.333 0.07006207
[4,] 48.33333 0.27106362 1951.333 0.9209689 1864.667 0.08904198

> plot( mw )
```

Other models II

```
> ##
> my <- apc.fit( A = lungDK$Ax,
+               P = lungDK$Px,
+               D = lungDK$D,
+               Y = lungDK$Y/10^5,
+               npar = 8,
+               ref.c = 1900,
+               dr.extr = "y" ) # person-yras, weight Y

NOTE: npar is specified as:
A P C
8 8 8
[1] "ML of APC-model Poisson with log(Y) offset : ( ACP ):\n"
      Model Mod. df. Mod. dev. Test df. Test dev. Pr(>Chi) Test dev/di
1      Age      212 15468.603      NA      NA      NA      NA
2      Age-drift 211 6858.883      1 8609.7199 0.00000e+00 8609.7199
3      Age-Cohort 205 1034.737      6 5824.1456 0.00000e+00 970.6909
4 Age-Period-Cohort 199 423.158      6 611.5791 7.41219e-129 101.9298
```

How to? III

```
cp.offset RR.fac
1765 1

> mw$Drift
exp(Est.) 2.5% 97.5%
APC (Y-weights) 1.020305 1.019450 1.021161
A-d 1.023487 1.022971 1.024003
```

Other models III

```
5      Age-Period 205 3082.602      6 2659.4439 0.00000e+00 443.2407
6      Age-drift 211 6858.883      6 3776.2808 0.00000e+00 629.3801

> ##
> ml <- apc.fit( A = lungDK$Ax,
+               P = lungDK$Px,
+               D = lungDK$D,
+               Y = lungDK$Y/10^5,
+               npar = 8,
+               ref.c = 1900,
+               dr.extr = "1" ) # usual inner product
```

Other models IV

```
NOTE: npar is specified as:
A P C
8 8 8
[1] "ML of APC-model Poisson with log(Y) offset : ( APC ):\n"
      Model Mod. df. Mod. dev. Test df. Test dev. Pr(>Chi) Test dev/df
1      Age      212 15468.603      NA      NA      0.00000e+00 8609.7199
2      Age-drift 211 6858.883      1 8609.7199 0.00000e+00 8609.7199
3      Age-Cohort 205 1034.737      6 5824.1456 0.00000e+00 970.6903
4 Age-Period-Cohort 199 423.158      6 611.5791 7.41219e-129 101.9298
5      Age-Period 205 3082.602      6 2659.4439 0.00000e+00 443.2407
6      Age-drift 211 6858.883      6 3776.2808 0.00000e+00 629.3803

> ##
> dr <- cbind( mw$Drift, ml$Drift, my$Drift, m1$Drift )
> rownames(dr) <- c("APC extract", "Age-Drift")
> colnames(dr)[0:3*3+1] <- c("D-wt", "Y^2/D-wt", "Y-wt", "1-wt")
> round( dr, 2 )
```

APC-model: Parametrization (APC-par)

252/ 333

Lee-Carter model for (mortality) rates

Lee & Carter, JASA, 1992:

$$\log(\lambda_{x,t}) = a_x + b_x \times k_t$$

x is age; t is calendar time

- ▶ Formulated originally using as step-functions with one parameter per age/period.
- ▶ Implicitly assumes a data lay out by age and period: A, B or C-sets, but **not** Lexis triangles
- ▶ Using Lexis triangles with categorical set-up would just produce separate models for upper and lower triangles.

Lee-Carter model (LeeCarter)

256/ 333

Other models V

```
D-wt 2.5% 97.5% Y^2/D-wt 2.5% 97.5% Y-wt 2.5% 97.5% 1-wt 2.5% 97.5%
APC extract 1.02 1.02 1.02 1.02 1.02 1.02 1.02 1.02 1.02 1.03 1.03 1.03
Age-Drift 1.02 1.02 1.02 1.02 1.02 1.02 1.02 1.02 1.02 1.02 1.02 1.02

> # % change per year
> round( (dr-1)*100, 1 )

D-wt 2.5% 97.5% Y^2/D-wt 2.5% 97.5% Y-wt 2.5% 97.5% 1-wt 2.5% 97.5%
APC extract 2.0 1.9 2.1 2.0 1.9 2.1 2.0 1.9 2.1 3.3 3.2 3.4
Age-Drift 2.3 2.3 2.4 2.3 2.3 2.4 2.3 2.3 2.4 2.3 2.3 2.4
```

Substantial differences between the estimated drifts.

APC-model: Parametrization (APC-par)

253/ 333

Lee-Carter model in continuous time

For **any** set of subsets of a Lexis diagram:

$$\log(\lambda(a, t)) = f(a) + b(a) \times k(t)$$

- ▶ $f(a)$, $b(a)$ smooth functions of age, a is **quantitative**
- ▶ $k(t)$ smooth function of period, t is **quantitative**
- ▶ Relative **scaling** of $b(a)$ and $k(t)$ cannot be determined
- ▶ $k(t)$ only determined up to an **affine** transformation:

$$\begin{aligned} f(a) + b(a)k(t) &= f(a) + (b(a)/n)(m + k(t) \times n) \\ &\quad - (b(a)/n) \times m \\ &= \tilde{f}(a) + \tilde{b}(a)\tilde{k}(t) \end{aligned}$$

Lee-Carter model (LeeCarter)

257/ 333

Lee-Carter model in continuous time

$$\log(\lambda(a, t)) = f(a) + b(a) \times k(t)$$

- ▶ Lee-Carter model is an extension of the age-period model; if $b(a) = 1$ it is the age-period model.
- ▶ The extension is an age \times period interaction, but not a traditional one:

$$\log(\lambda(a, t)) = f(a) + b(a) \times k(t) = f(a) + k(t) + (b(a) - 1) \times k(t)$$

- ▶ Main effect and interaction component of t are constrained to be identical.

Lee-Carter model (LeeCarter)

258/ 333

Parametrization of the APC model is arbitrary

- ▶ Separation of the three effects relies on arbitrary principles, e.g.:
 - ▶ Age is the primary effect
 - ▶ Cohort the secondary, reference c_0
 - ▶ Period is the residual
 - ▶ Inner product for trend extraction
- ▶ There is no magical fix that allows you to escape this, it comes from modelling a , p and $p - a$
- ▶ Any fix has some (hidden) assumption(s)
- ▶ ... but the **fitted values** are the same

APC-model: Parametrization (APC-par)

255/ 333

Main effect and interaction term

Main effect and interaction component of t are constrained to be identical.

None of these are Lee-Carter models:

```
> glm( D ~ Ns(A, kn=a1.kn) + Ns(A, kn=a2.kn, i=T) : Ns(P, kn=p.kn), ... )
> glm( D ~ Ns(A, kn=a1.kn) + Ns(A, kn=a2.kn, i=T) * Ns(P, kn=p.kn), ... )
> glm( D ~ Ns(A, kn=a1.kn) + Ns(P, kn=p.kn) + Ns(A, kn=a2.kn, i=T) : Ns(P, kn=p.kn), ... )
```

Lee-Carter model (LeeCarter)

259/ 333

Lee-Carter model

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

LeeCarter

Lee-Carter model interpretation

$$\log(\lambda(a, p)) = f(a) + b(a) \times k(p)$$

- ▶ Constraints:
- ▶ $f(a)$ is the basic age-specific mortality
- ▶ $k(p)$ is the rate-ratio (RR) as a function of p :
 - ▶ relative to a p_{ref} where $k(p_{ref}) = 1$
 - ▶ for persons aged a_{ref} where $b(a_{ref}) = 1$
- ▶ $b(a)$ is an age-specific multiplier for the RR $k(p)$
- ▶ Choose p_{ref} and a_{ref} a priori.

Lee-Carter model (LeeCarter)

260/ 333

Danish lung cancer data I

```
> lung <- read.table("../data/apc-Lung.txt", header=T)
> head(lung)
  sex A   P   C D   Y
1  1 0 1943 1942 0 19546.2
2  1 0 1943 1943 0 20796.5
3  1 0 1944 1943 0 20681.3
4  1 0 1944 1944 0 22478.5
5  1 0 1945 1944 0 22369.2
6  1 0 1945 1945 0 23885.0

> # Only A by P classification - and only men over 40
> ltab <- xtabs(cbind(D,Y) ~ A + P, data=subset(lung,sex==1))
> str(ltab)
```

Lee-Carter re-scaled I

$$\log(\hat{\lambda}(a, p)) = [f(a) + b(a) \times 20] + [b(a) \times 50] \times [(k(t) - 20)/50]$$

```
> par(mfcol=c(1,3))
> matplot(dmg.lcM$age, exp(dmg.lcM$ax+dmg.lcM$bx*20)*1000,
+         log="y", ylab="Lung cancer incidence rates per 1000 PY",
+         xlab="Age", type="l", lty=1, lwd=4)
> matplot(dmg.lcM$age, dmg.lcM$bx*50,
+         ylab="Age effect",
+         xlab="Age", type="l", lty=1, lwd=4)
> abline(h=1)
> matplot(dmg.lcM$year, (dmg.lcM$kt-20)/50,
+         ylab="Time effect",
+         xlab="Date", type="l", lty=1, lwd=4)
> abline(h=0)
```


Danish lung cancer data II

```
xtabs [1:90, 1:61, 1:2] 0 0 0 0 0 0 0 0 0 0 ...
- attr(*, "dimnames")=List of 3
..$ A: chr [1:90] "0" "1" "2" "3" ...
..$ P: chr [1:61] "1943" "1944" "1945" "1946" ...
..$ : chr [1:2] "D" "Y"
- attr(*, "call")= language xtabs(formula = cbind(D, Y) ~ A + P, data = subset(lung, sex == 1))
```

Lee-Carter modeling in R-packages:

- ▶ demography (`lca`)
- ▶ ilc (`lca.rh`)
- ▶ Epi (`LCa.fit`).

Lee-Carter with demography rescaled



Lee-Carter with demography I

```
> library(demography)
> lcM <- demogdata(data = as.matrix(ltab[40:90, "D"]/ltab[40:90, "Y"]),
+                 pop = as.matrix(ltab[40:90, "Y"]),
+                 ages = as.numeric(dimnames(ltab)[1:1][40:90]),
+                 years = as.numeric(dimnames(ltab)[2:2]),
+                 type = "Lung cancer incidence",
+                 label = "Denmark",
+                 name = "Male")
```

`lca` estimation function checks the `type` argument, so we make a work-around, `mrt`:

Lee-Carter with ilc

- ▶ The `lca.rh` function fits the model using maximum likelihood (proportional scaling)
- ▶ Fits the more general model and submodels of it:

$$\log(\lambda(a, p)) = f(a) + b(a) \times k(p) + c(a)m(p - a)$$

- ▶ Age interaction with between age and both period and/or cohort (=period-age)
 - ▶ It is also an extension of the APC-model; if $b(a) = 1$ and $c(a) = 1$ it's the APC-model.
- ⇒ suffers from the same identifiability problem

Lee-Carter with demography II

```
> mrt <- function(x) { x$type <- "mortality" ; x }
> dmg.lcM <- lca(mrt(lcM), interpolate=TRUE)
> par(mfcol=c(1,3))
> matplot(dmg.lcM$age, exp(dmg.lcM$ax)*1000,
+         log="y", ylab="Lung cancer incidence rates per 1000 PY",
+         xlab="Age", type="l", lty=1, lwd=4)
> matplot(dmg.lcM$age, dmg.lcM$bx,
+         ylab="Age effect",
+         xlab="Age", type="l", lty=1, lwd=4)
> matplot(dmg.lcM$year, dmg.lcM$kt,
+         ylab="Time effect",
+         xlab="Date", type="l", lty=1, lwd=4)
> abline(h=0)
```

Lee-Carter with ilc I

```
> library(ilc)
> ilc.lcM <- lca.rh(mrt(lcM), model="lc", interpolate=TRUE, verbose=FALSE)
```

Original sample: Mortality data for Denmark
Series: Male
Years: 1943 - 2003
Ages: 39 - 89


Applied sample: Mortality data for Denmark (Corrected: interpolate)
Series: Male
Years: 1943 - 2003
Ages: 39 - 89

Fitting model: [LC = a(x)+b1(x)*k(t)]
- with poisson error structure and with deaths as weights -


Iterations finished in: 34 steps

```
> plot(ilc.lcM)
```

Lee-Carter with demography



Lee-Carter with ilc



Lee-Carter with Epi

- ▶ `LCa.fit` fits the Lee-Carter model using natural splines for the **quantitative** effects of age and time.
- ▶ Normalizes effects to a reference age and period.
- ▶ The algorithm alternately fits a main age and period effects and the age-interaction effect.


$$\log(\lambda(a, p)) = f(a) + b(a) \times k(p) + c(a) \times m(p - a)$$

$$\log(\lambda(a, p)) = f(a) + b(a) \times k(p) + c(a) \times m(p - a)$$

Lee-Carter model (LeeCarter)

271/ 333

Lee-Carter models and APC models



Lee-Carter model (LeeCarter)

276/ 333

Lee-Carter with Epi I

```
> library( Epi )
> M1c <- subset( lung, sex==1 & A>39 )
> LCa.M1c <- LCa.fit( M1c, a.ref=60, p.ref=1980 )

LCa.fit convergence in 8 iterations, deviance: 8548.443 on 6084 d.f.
> LCa.M1c

APa: Lee-Carter model with natural splines:
log(Rate) = ax(Age) + pi(Age)kp(Per)
with 6, 5 and 5 parameters respectively.
Deviance: 8548.443 on 6084 d.f.

> plot( LCa.M1c, rnam="Lung cancer incidence per 1000 PY" )
```

Lee-Carter model (LeeCarter)

272/ 333


Lee-Carter models and APC models

- ▶ The classical Lee-Carter model is an extension of the Age-Period model with an interaction
- ▶ The Age-Period-Cohort model is an extension of the Age-Period model with an interaction
- ▶ Replacing period with cohort gives another type of Lee-Carter model
- ▶ The logical step is to consider all 9 models that comes from cross-classification of how the interaction term $b(a)$
 - ▶ Linear effect ($b(a) = 0$)
 - ▶ Non-linear effect ($b(a) = 1$)
 - ▶ Multiplicative interaction with age ($b(a)$ unconstrained)

Lee-Carter model (LeeCarter)

277/ 333

Lee-Carter with Epi



Lee-Carter model (LeeCarter)

273/ 333

Lee-Carter models and APC models

		$b_c(a)$		
		0	1	free
$b_p(a)$	0	Age	Age+Coh	LCa(C) <i>AC, ac, ACa</i>
	1	Age+Per	Age+Per+Coh <i>H₀, h₀</i>	Age+Per+LCa(C) <i>H₁, h₁, APCa</i>
	free	LCa(P) <i>LC, lc, APa</i>	Age+Coh+LCa(P) <i>H₂, h₂, APaC</i>	Age+LCa(P)+LCa(C) <i>M, m, APaCa</i>

Model: `ilc: lca.rh(model=)` Epi: `LCa.fit(model=)`

Lee-Carter model (LeeCarter)

278/ 333

Lee-Carter and the APC-model

- ▶ Lee-Carter model is an interaction extension of the Age-Period model
- ▶ ... or an interaction extension of the Age-Cohort model
- ▶ Age-Period-Cohort model is:
 - ▶ interaction extension
 - ▶ the smallest **union** of Age-Period and Age-Cohort
- ▶ Extended Lee-Carter (from the `ilc` package)

$$\log(\lambda(a, p)) = f(a) + b(a) \times k(p) + c(a)m(p - a)$$

is the union of all of these.

Lee-Carter model (LeeCarter)

274/ 333

Wednesday afternoon exercise: Age-period-cohort model

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

exc-Wed2

Lee-Carter and the APC-model

```
> system.time( allmod <- apc.LCa( M1c, keep.models=TRUE ) )
> str( allmod )
> save( allmod, file='allmod.Rda' )

> load( file='allmod.Rda' )
> show.apc.LCa( allmod, top="Ad" )

> show.apc.LCa( allmod, top="AP" )

> show.apc.LCa( allmod, top="AC" )
```

Lee-Carter model (LeeCarter)

275/ 333

Wednesday exercise

1. Load the `Epi` package.
2. Do the practical 2.5: Age-period-cohort models
3. I will be available the first approx. hour for direct questions that will be broadcast to the entire audience.
4. After that I will be on e-mail, until about 15.
5. Thursday morning we will:
 - ▶ Recap the contents of today's lecture
 - ▶ Go over the solutions to today's exercises
6. Remember: Questions always welcome at any time, just switch on you microphone and peak.

Wednesday afternoon exercise: Age-period-cohort model (exc-Wed2)

279/ 333

APC-models for several datasets

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:

Age-Period-Cohort models

— and some cousins

European Doctoral School of Demography,

Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

APC2

```
> apc.Sem <- apc.fit( subset( th, hist=="Sem" ),
+                   parm = "ACP",
+                   ref.c = 1970,
+                   npar = c(A=8,P=8,C=8) )

[1] "ML of APC-model Poisson with log(Y) offset : ( ACP ):\n"
      Model Mod. df. Mod. dev. Test df. Test dev. Pr(>Chi) Test dev/c
1      Age      5392  5202.544      NA      NA      NA      NA
2 Age-drift    5391  5074.144      1 603.33315 3.153666e-133 603.33315
3 Age-Cohort   5385  5038.675      6 35.46902 3.495353e-06 5.91150
4 Age-Period-Cohort 5379 5014.665      6 24.00981 5.200936e-04 4.00163
5 Age-Period   5385 5061.467      6 46.80254 2.048715e-08 7.80042
6 Age-drift    5391  5074.144      6 12.67628 4.847449e-02 2.11272

> apc.nS <- apc.fit( subset( th, hist=="nS" ),
+                   parm = "ACP",
+                   ref.c = 1970,
+                   npar = c(A=8,P=8,C=8) )
```

APC-models for several datasets (APC2)

284/ 333

Two APC-models

- APC-models for two sets of rates (men/women, say)

$$\log(\lambda_i(a, p)) = f_i(a) + g_i(p) + h_i(p - a), \quad i = 1, 2$$

- Rate-ratio also an APC-model:

$$\begin{aligned} \log(\text{RR}(a, p)) &= \log(\lambda_1(a, p)) - \log(\lambda_2(a, p)) \\ &= (f_1(a) - f_2(a)) + (g_1(p) - g_2(p)) \\ &\quad + (h_1(p - a) - h_2(p - a)) \\ &= f_{\text{RR}}(a) + g_{\text{RR}}(p) + h_{\text{RR}}(p - a) \end{aligned}$$

- Model the two sets of rates separately and report the ratio effects as any other APC-model.
- Note; not all constraints carry over to RR

APC-models for several datasets (APC2)

280/ 333

```
[1] "ML of APC-model Poisson with log(Y) offset : ( ACP ):\n"
      Model Mod. df. Mod. dev. Test df. Test dev. Pr(>Chi) Test dev/c
1      Age      5392  5202.544      NA      NA      NA      NA
2 Age-drift    5391  4501.466      1 701.07777 1.743153e-154 701.07777
3 Age-Cohort   5385  4459.701      6 41.76543 2.045644e-07 6.96090
4 Age-Period-Cohort 5379 4375.172      6 84.52883 4.132959e-16 14.08813
5 Age-Period   5385  4427.599      6 52.42632 1.530676e-09 8.73772
6 Age-drift    5391  4501.466      6 73.86794 6.563086e-14 12.31132

> round( cbind( apc.Sem$Drift,
+              apc.nS$Drift ) -1)*100, 1 )

      exp(Est.) 2.5% 97.5% exp(Est.) 2.5% 97.5%
APC (Y-weights) 2.6 2.4 2.9      3.4 3.0 3.7
A-d             2.5 2.3 2.7      3.1 2.8 3.3

> plot( apc.Sem, "Sem vs. non-Sem RR", col="transparent" )

cp.offset  RR.fac
1804       1
```

APC-models for several datasets (APC2)

285/ 333

Two sets of data I

Example: Testis cancer in Denmark, Seminoma and non-Seminoma cases.

```
> th <- read.table( "../data/testis-hist.txt", header=TRUE )
> str( th )

'data.frame': 29160 obs. of 9 variables:
 $ a : int 0 0 0 0 0 1 1 1 1 ...
 $ p : int 1943 1943 1943 1943 1943 1943 1943 1943 1943 ...
 $ c : int 1942 1942 1942 1943 1943 1943 1941 1941 1942 ...
 $ y : num 18853 18853 18853 20796 20796 ...
 $ age : num 0.667 0.667 0.667 0.333 0.333 ...
 $ diag : num 1943 1943 1943 1944 1944 ...
 $ birth: num 1943 1943 1943 1943 1943 ...
 $ hist : int 1 2 3 1 2 3 1 2 3 1 ...
 $ d : int 0 1 0 0 0 0 0 0 0 ...
```

APC-models for several datasets (APC2)

281/ 333

```
> matshade( apc.nS$Age[,1], ci.ratio( apc.Sem$Age[, -1], apc.nS$Age[, -1] ), col=1 )
> pc.matshade( apc.nS$Per[,1], ci.ratio( apc.Sem$Per[, -1], apc.nS$Per[, -1] ), col=1 )
> pc.matshade( apc.nS$Coh[,1], ci.ratio( apc.Sem$Coh[, -1], apc.nS$Coh[, -1] ), col=1 )
> abline( h=1 )
```

APC-models for several datasets (APC2)

286/ 333

Two sets of data II


```
> head( th )

  a  p  c  y  age  diag  birth hist d
1 0 1943 1942 18853.0 0.6666667 1943.333 1942.667 1 0
2 0 1943 1942 18853.0 0.6666667 1943.333 1942.667 2 1
3 0 1943 1942 18853.0 0.6666667 1943.333 1942.667 3 0
4 0 1943 1943 20796.5 0.3333333 1943.667 1943.333 1 0
5 0 1943 1943 20796.5 0.3333333 1943.667 1943.333 2 0
6 0 1943 1943 20796.5 0.3333333 1943.667 1943.333 3 0

> th <- transform( th,
+                 hist = factor( hist, labels=c("Sem", "nS", "Oth" ) ),
+                 A = age,
+                 P = diag,
+                 D = d,
+                 Y = y/10^4 ), c("A", "P", "D", "Y", "hist" )
> th <- subset( th, A>15 & A<65 & hist!="Oth" )
> th$hist <- factor( th$hist )
```

APC-models for several datasets (APC2)

282/ 333



APC-models for several datasets (APC2)

287/ 333


```
> library( Epi )
> stat.table( list( Histology = hist ),
+            list( D = sum(D),
+                  Y = sum(Y) ),
+            margins = TRUE,
+            data = th )
```

Histology	D	Y
Sem	4461.00	8435.49
nS	3494.00	8435.49
Total	7955.00	16870.99

First step is separate analyses for each subtype (Sem, nS, resp.)

APC-models for several datasets (APC2)

283/ 333



APC-models for several datasets (APC2)

287/ 333

Analysis of two rates: Formal tests I

Separate models with the same parametrization:

```
> ( Akn <- (apc.sem$Knots$Age+apc.ns$Knots$Age)/2 )
[1] 22.66667 26.50000 29.50000 32.33333 35.16667 38.83333 43.83333 52.66667
> ( Pkn <- (apc.sem$Knots$Per+apc.ns$Knots$Per)/2 )
[1] 1952.417 1964.000 1972.333 1978.167 1983.000 1987.500 1991.500 1995.000
> ( Ckn <- (apc.sem$Knots$Coh+apc.ns$Knots$Coh)/2 )
[1] 1913.500 1926.000 1934.833 1942.000 1947.833 1953.333 1958.958 1966.000
> apc.sem <- apc.fit( subset(th,hist=="Sem"), npar=list(A=Akn,P=Pkn,C=Ckn), pr=F )
No reference cohort given; reference cohort for age-effects is chosen as
the median date of birth for persons with event: 1939.667 .
```

APC-model: Interactions

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

Analysis of two rates: Formal tests II

```
> apc.ns <- apc.fit( subset(th,hist=="nS"), npar=list(A=Akn,P=Pkn,C=Ckn), pr=F )
No reference cohort given; reference cohort for age-effects is chosen as
the median date of birth for persons with event: 1949.667 .
```

Joint model, parametrize interactions separately:

```
> Ma <- with( th, Ns( A, knots=Akn, intercept=TRUE ) )
> Mp <- with( th, Ns( P, knots=Pkn ) )
> Mc <- with( th, Ns( P-A, knots=Ckn ) )
> # extract the linear trend
> Mp <- detrend( Mp, th$P, weight=th$D )
> Mc <- detrend( Mc, th$P-th$A, weight=th$D )
> m.apc <- glm( D ~ -1 + Ma:hist + Mp:hist + Mc:hist +
+ P:hist + # note separate slopes extracted
+ offset( log(Y) ),
+ family=poisson, data=th )
> m.apc$deviance
```

Analysis of DM-rates: Age×sex interaction I

- ▶ 10 centres
- ▶ 2 sexes
- ▶ Age: 0–15
- ▶ Period 1989–1999
- ▶ Is the sex-effect the same between all centres?
- ▶ How is timetrend by birth cohort?

Analysis of two rates: Formal tests III

```
[1] 9410.446
> # Same as the sum from separate models
> apc.ns$Model$deviance + apc.sem$Model$deviance
[1] 9410.446
```

Tests for equality of non-linear part of shapes

```
> m.ap <- update( m.apc, . ~ . - Mc:hist + Mc )
> m.ac <- update( m.apc, . ~ . - Mp:hist + Mp )
> m.a <- update( m.ap, . ~ . - Mp:hist + Mp )
> m.d <- update( m.ap, . ~ . - Mp:hist )
> m.o <- update( m.ap, . ~ . - P:hist + P )
> AOV <- anova( m.a, m.ac, m.apc, m.ap, m.a, m.d, m.o, test="Chisq")
> rownames( AOV ) <- c("","cohRR","perRR|coh","cohRR|per","perRR","drift","Smdrift")
> AOV
```

Analysis of DM-rates: Age×sex interaction II

```
> library( Epi )
> library( splines )
> # load( file="c:/Bendix/Artikler/A_P_C/IDDM/Eurodiab/data/tri.Rdata" )
> load( file = "~/teach/APC/examples/EuroDiab/tri.Rdata" )
> str(dm)

'data.frame': 5940 obs. of 8 variables:
 $ sex: Factor w/ 2 levels "F","M": 1 1 1 1 1 1 1 1 1 1 ...
 $ cen: Factor w/ 10 levels "Z2: Czech","A1: Austria",...: 2 2 2 2 2 2 2 2 2 2 ...
 $ per: num 1989 1990 1991 1992 1993 ...
 $ D : num 1 0 0 0 0 0 0 0 1 ...
 $ A : num 0.333 0.333 0.333 0.333 0.333 0.333 ...
 $ P : num 1990 1991 1992 1993 1994 ...
 $ C : num 1989 1990 1991 1992 1993 ...
 $ Y : num 21970 22740 22886 23026 22323 ...
```

Analysis of two rates: Formal tests IV

Analysis of Deviance Table

```
Model 1: D ~ Mc + Mp + Ma:hist + hist:P + offset(log(Y)) - 1
Model 2: D ~ Mp + Ma:hist + hist:Mc + hist:P + offset(log(Y)) - 1
Model 3: D ~ -1 + Ma:hist + Mp:hist + Mc:hist + P:hist + offset(log(Y))
Model 4: D ~ Mc + Ma:hist + hist:Mp + hist:P + offset(log(Y)) - 1
Model 5: D ~ Mc + Mp + Ma:hist + hist:P + offset(log(Y)) - 1
Model 6: D ~ Mc + Ma:hist + hist:P + offset(log(Y)) - 1
Model 7: D ~ Mc + P + Ma:hist + hist:Mp + offset(log(Y)) - 1
  Resid. Df Resid. Dev Df Deviance Pr(>Chi)
cohRR      10770      9467.4
perRR|coh  10764      9447.3 6 20.094 0.002665
cohRR|per  10758      9410.4 6 36.886 1.854e-06
perRR      10764      9421.6 -6 -11.196 0.082496
drift      10770      9467.4 -6 -45.783 3.270e-08
Smdrift    10776      9538.2 -6 -70.807 2.793e-13
```

Analysis of DM-rates: Age×sex interaction III

```
> dm <- dm[dm$cen=="D1: Denmark",]
> attach( dm )
> # Define knots and points of prediction
> n.A <- 5
> n.C <- 8
> n.P <- 5
> c0 <- 1985
> attach( dm, warn.conflicts=FALSE )
> A.kn <- quantile( rep( A, D ), probs=(1:n.A-0.5)/n.A )
> P.kn <- quantile( rep( P, D ), probs=(1:n.P-0.5)/n.P )
> C.kn <- quantile( rep( C, D ), probs=(1:n.C-0.5)/n.C )
> A.pt <- sort( A[match( unique(A), A )] )
> P.pt <- sort( P[match( unique(P), P )] )
> C.pt <- sort( C[match( unique(C), C )] )
> # Age-cohort model with age-sex interaction
> # The model matrices for the ML fit
> # - note that intercept is in age term, and drift is added to the cohort term:
> Ma <- Ns( A, kn=A.kn, intercept=T )
> Mc <- cbind( C-c0, detrend( Ns( C, kn=C.kn ), C, weight=D ) )
```

Several datasets I

- ▶ Separate models for each
- ▶ Rate-ratios between two sets of fitted rates also follow an APC model
- ▶ Constraints does not necessarily carry over to RRs
- ▶ Test for equality of effects: non-linear and linear
- ▶ Take care not to violate the **principle of marginality**: — do not test linear terms when non-linear terms are in the model.

Analysis of DM-rates: Age×sex interaction IV

```
> Mp <- detrend( Ns( P, kn=P.kn ), P, weight=D )
> # The prediction matrices - corresponding to ordered unique values of A, P and C
> Pa <- Ma[match(A.pt,A),,drop=F]
> Pp <- Mp[match(P.pt,P),,drop=F]
> Pc <- Mc[match(C.pt,C),,drop=F]
> # Fit the apc model using the cohort major parametrization
> apcs <- glm( D ~ Ma:sex - 1 + Mc + Mp +
+ offset( log( Y/10^5 ) ),
+ family=poisson, epsilon = 1e-10,
+ data=dm )
> ci.exp( apcs )
```

Analysis of DM-rates: Age×sex interaction V

```

exp(Est.) 2.5% 97.5%
Mc 1.0053157 0.9719640 1.0398118
Mc1 0.6496197 0.3305926 1.2765132
Mc2 1.2576228 0.6868926 2.3025652
Mc3 0.5366885 0.2787860 1.0331743
Mc4 0.9207689 0.4877809 1.7381069
Mc5 0.6898805 0.3999550 1.1899714
Mc6 1.1005438 0.5817089 2.0821352
Mp1 0.5735223 0.3489977 0.9424928
Mp2 1.0534148 0.6090201 1.8220792
Mp3 0.9412582 0.4032633 2.1969937
Ma1:sexF 11.9104421 6.7605869 20.9831831
Ma2:sexF 22.0985163 11.9531639 40.8548253
Ma3:sexF 16.5201055 9.6623215 28.2451673
Ma4:sexF 360.8119685 225.4568974 577.4286708
Ma5:sexF 2.5694234 1.5219041 4.3379452
Ma1:sexM 17.0238730 9.9414867 29.1518021
Ma2:sexM 13.4664178 7.0861312 25.5914549
Ma3:sexM 14.4664367 8.6164003 24.2883087

```

APC-model: Interactions (APC-Int)

297 / 333

Analysis of DM-rates: Age×sex interaction X

```

> matshade( A.pt, M.inc, lwd=2, col="blue" )
> matshade( A.pt, F.inc, lwd=2, col="red" )
> matshade( A.pt, MF.RR*5, lwd=2 ) ; abline( h=5 )
> pc.matshade( C.pt, c.RR, lwd=2 )
> pc.matshade( P.pt, p.RR, lwd=2 )

```

APC-model: Interactions (APC-Int)

302 / 333

Analysis of DM-rates: Age×sex interaction VI

```

Ma4:sexM 531.9214375 343.2221445 824.3652694
Ma5:sexM 3.1485499 1.9406858 5.1081770

```

```

> # Average trend (D-projection)
> round( ( ci.exp( apcs, subset=1 ) - 1 ) *100, 1 )

exp(Est.) 2.5% 97.5%
Mc 0.5 -2.8 4

> ci.exp( apcs, subset="sexF" )


exp(Est.) 2.5% 97.5%
Ma1:sexF 11.910442 6.760587 20.983183
Ma2:sexF 22.098516 11.953164 40.854825
Ma3:sexF 16.520106 9.662321 28.245167
Ma4:sexF 360.811968 225.456897 577.428671
Ma5:sexF 2.569423 1.521904 4.337945

> cbind( A.pt, ci.exp( apcs, subset="sexF", ctr.mat=Pa ) )

```

APC-model: Interactions (APC-Int)

298 / 333



APC-model: Interactions (APC-Int)

303 / 333

Analysis of DM-rates: Age×sex interaction VII

```

A.pt exp(Est.) 2.5% 97.5%
[1,] 0.3333333 4.943285 2.363023 10.34102
[2,] 0.6666667 5.309563 2.676029 10.53481
[3,] 1.3333333 6.125551 3.416160 10.98379
[4,] 1.6666667 6.579431 3.847562 11.25100
[5,] 2.3333333 7.590575 4.833655 11.91993
[6,] 2.6666667 8.153008 5.380890 12.35326
[7,] 3.3333333 9.401089 6.531373 13.53168
[8,] 3.6666667 10.085197 7.103019 14.31943
[9,] 4.3333333 11.561158 8.190634 16.31869
[10,] 4.6666667 12.344483 8.712446 17.49064
[11,] 5.3333333 13.969938 9.777715 19.95959
[12,] 5.6666667 14.794673 10.355375 21.13708
[13,] 6.3333333 16.412682 11.674678 23.07354
[14,] 6.6666667 17.179232 12.425801 23.75107
[15,] 7.3333333 18.578132 13.958075 24.72741
[16,] 7.6666667 19.228123 14.579373 25.35917
[17,] 8.3333333 20.513353 15.309646 27.48579
[18,] 8.6666667 21.190703 15.538953 28.89808

```

APC-model: Interactions (APC-Int)

299 / 333

Analysis of DM-rates: Age×sex interaction I

A bit more intuitive, independent of parametrization:

```

> apcS <- glm( D ~ Ns(A,knots=A.kn,intercept=TRUE):sex +
+ Ns(P,knots=P.kn) + Ns(C,knots=C.kn) +
+ offset( log( Y/10^5 ) ),
+ family=poisson, epsilon = 1e-10,
+ data=dm )
> apcS$deviance
[1] 633.5838

> apcs$deviance
[1] 633.5838

```

APC-model: Interactions (APC-Int)

304 / 333

Analysis of DM-rates: Age×sex interaction VIII

```

[19,] 9.3333333 22.742587 16.317839 31.69692
[20,] 9.6666667 23.679333 17.100960 32.78827
[21,] 10.3333333 25.893547 19.499950 34.38346
[22,] 10.6666667 26.999519 20.607727 35.37382
[23,] 11.3333333 28.605296 21.348779 38.32832
[24,] 11.6666667 28.831963 21.013988 39.55851
[25,] 12.3333333 27.526786 19.701501 38.46022
[26,] 12.6666667 25.941507 18.827598 35.74337
[27,] 13.3333333 21.900696 16.035816 29.91058
[28,] 13.6666667 19.869417 14.038380 28.12246
[29,] 14.3333333 16.320075 10.026866 26.56312
[30,] 14.6666667 14.790766 8.323640 26.28258

> # Extract the effects
> F.inc <- ci.exp( apcs, subset="sexF", ctr.mat=Pa)
> M.inc <- ci.exp( apcs, subset="sexM", ctr.mat=Pa)
> MF.RR <- ci.exp( apcs, subset=c("sexM","sexF"), ctr.mat=cbind(Pa,-Pa))
> c.RR <- ci.exp( apcs, subset="Mc", ctr.mat=Pc)
> p.RR <- ci.exp( apcs, subset="Mp", ctr.mat=Pp)

```

APC-model: Interactions (APC-Int)

300 / 333

Analysis of DM-rates: Age×sex interaction II

```

> # rates for the 1985 birth cohort and the RR
> a.pt <- seq(0,15,0.1)
> ndaM <- data.frame( A=a.pt, P=1985+a.pt, C=1985, Y=10^5, sex="M" )
> ndaF <- data.frame( A=a.pt, P=1985+a.pt, C=1985, Y=10^5, sex="F" )
> a.pM <- ci.pred( apcS, ndaM )
> a.pF <- ci.pred( apcS, ndaF )
> a.RR <- ci.exp( apcs, list(ndaM,ndaF) )
> # Cohort RRs relative to C=1985
> ndc <- data.frame( A=10, P=2000, C=1975:2000, Y=10^5 )
> ndr <- data.frame( A=10, P=2000, C=1985, Y=10^5 )
> c.RR <- ci.exp( apcS, list(ndc,ndr) )
> # Period RRs relative to P=2000
> ndp <- data.frame( A=10, P=1990:2000, C=1985, Y=10^5 )
> ndr <- data.frame( A=10, P=2000, C=1985, Y=10^5 )
> p.RR <- ci.exp( apcS, list(ndp,ndr) )
> # plt( paste( "DM-DK" ), width=11 )
> par( mar=c(4,4,1,4), mgp=c(3,1,0)/1.6, las=1 )
> #
> # The the frame for the effects

```

APC-model: Interactions (APC-Int)

305 / 333

Analysis of DM-rates: Age×sex interaction IX

The the frame for the effects

```

> par( mar=c(4,4,1,4), mgp=c(3,1,0)/1.6, las=1 )
> apc.frame( a.lab=c(0,5,10,15),
+ a.tic=c(0,5,10,15),
+ r.lab=c(c(1,1.5,3,5),c(1,1.5,3,5)*10),
+ r.tic=c(c(1,1.5,2,5),c(1,1.5,2,5)*10),
+ cp.lab=seq(1980,2000,10),
+ cp.tic=seq(1975,2000,5),
+ rr.ref=5,
+ gap=1,
+ col.grid=gray(0.9),
+ a.txt="",
+ cp.txt="",
+ r.txt="",
+ rr.txt="" )
> # Draw the estimates
> matshade( a.pt, a.pM, lwd=2, col="blue" )
> matshade( a.pt, a.pF, lwd=2, col="red" )
> matshade( a.pt, a.RR*5, lwd=2 ) ; abline( h=5 )
> pc.matshade( 1975:2000, c.RR, lwd=2 )
> pc.matshade( 1990:2000, p.RR, lwd=2 )

```

APC-model: Interactions (APC-Int)

301 / 333

Analysis of DM-rates: Age×sex interaction III

```


> apc.frame( a.lab=c(0,5,10,15),
+ a.tic=c(0,5,10,15),
+ r.lab=c(c(1,1.5,3,5),c(1,1.5,3,5)*10),
+ r.tic=c(c(1,1.5,2,5),c(1,1.5,2,5)*10),
+ cp.lab=seq(1980,2000,10),
+ cp.tic=seq(1975,2000,5),
+ rr.ref=5,
+ gap=1,
+ col.grid=gray(0.9),
+ a.txt="",
+ cp.txt="",
+ r.txt="",
+ rr.txt="" )
> # Draw the estimates
> matshade( a.pt, a.pM, lwd=2, col="blue" )
> matshade( a.pt, a.pF, lwd=2, col="red" )
> matshade( a.pt, a.RR*5, lwd=2 ) ; abline( h=5 )
> pc.matshade( 1975:2000, c.RR, lwd=2 )
> pc.matshade( 1990:2000, p.RR, lwd=2 )

```




APC-model: Interactions (APC-Int)

306 / 333

Analysis of DM-rates: Age×sex interaction IV



- ▶ ... but these are not the estimates we really want as before.
- ▶ The detrended estimates are not available from the fitted values, because the parametrization they rely on is a function of **data**.
- ▶ Of course the parameters can be extracted but it requires a construction of the model matrices as we did first
- ▶ How is shown in the section "Reparametrizations" in the notes on "Introductory linear algebra with R".



Predicting future rates

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

predict

Prediction of future rates

Model:

$$\log(\lambda(a, p)) = f(a) + g(p) + h(c)$$

- ▶ Why not just extend the estimated functions into the future?
- ▶ Natural splines lend themselves easily to this [?]
- ▶ The parametrization curse — the model as stated is not uniquely parametrized.
- ▶ Predictions from the model must be invariant under reparametrization.

Identifiability

Predictions based in the three functions ($f(a)$, $g(p)$ and $h(c)$) must give the same prediction also for the reparametrized version:

$$\begin{aligned} \log(\lambda(a, p)) &= \tilde{f}(a) + \tilde{g}(p) + \tilde{h}(c) \\ &= (f(a) - \gamma a) + \\ &\quad (g(p) + \gamma p) + \\ &\quad (h(c) - \gamma c) \end{aligned}$$

A prediction based on the parametrization ($f(a)$, $g(p)$, $h(c)$) must give the same predictions as one based on ($\tilde{f}(a)$, $\tilde{g}(p)$, $\tilde{h}(c)$)

Parametrization invariance

- ▶ Prediction of the future course of g and h must preserve addition of a linear term in the argument:

$$\begin{aligned} \text{pred}(g(p) + \gamma p) &= \text{pred}(g(p)) + \gamma p \\ \text{pred}(h(c) - \gamma c) &= \text{pred}(h(c)) - \gamma c \end{aligned}$$

- ▶ If this is met, the predictions made will not depend on the parametrization chosen.
- ▶ If one of the conditions does not hold, the prediction will depend on the parametrization chosen.
- ▶ Any linear combination of (known) function values of $g(p)$ and $h(c)$ will work.


Identifiability

- ▶ Any linear combination of function values of $g(p)$ and $h(c)$ will work.
- ▶ Coefficients in the linear combinations used for g and h must be the same; otherwise the prediction will depend on the specific parametrization.
- ▶ What works best in reality is difficult to say: depends on the subject matter.

Predicting future rates (predict)

316/ 333

Example: Breast cancer in Denmark



Predicting future rates (predict)

317/ 333


Practicalities

- ▶ Long term predictions notoriously unstable.
- ▶ Decreasing slopes are possible, the requirement is that at any future point changes in the parametrization should cancel out in the predictions.

Predicting future rates (predict)

318/ 333

Breast cancer prediction



Predicted age-specific breast cancer rates at 2020 & 2030,

in the 1960 and 1970 cohorts.

Predicting future rates (predict)

319/ 333

APC-model for quantitative outcomes

- ▶ The classical model is:

$$\log(\lambda(a, p)) = f(a) + g(p) + h(p - a)$$

- ▶ In principle it would be possible to use an identity-link model:

$$\lambda(a, p) = f(a) + g(p) + h(p - a)$$

- ▶ ... or use APC-modelling for **measurement** data such as BMI, measured at different times and ages:

$$\text{BMI}_{ap} = f(a) + g(p) + h(p - a) + e_{ap}, \quad e_i \sim \mathcal{N}(0, \sigma^2)$$

- ▶ ... or more precisely:

$$\text{BMI}_i = f(a(i)) + g(p(i)) + h(p(i) - a(i)) + e_i, \quad e_i \sim \mathcal{N}(0, \sigma^2)$$

Continuous outcomes (cont)

320/ 333

APC-model for quantitative outcomes

- ▶ Model:

$$\text{BMI}_i = f(a(i)) + g(p(i)) + h(p(i) - a(i)) + e_i, \quad e_i \sim \mathcal{N}(0, \sigma^2)$$

- ▶ But the identification problem is still the same:

$$c(i) = p(i) - a(i), \quad \forall i$$


- ▶ But the same machinery applies with extraction of the effects
- ▶ — and plotting of predictions of
 - ▶ $E(\text{BMI})$
 - ▶ quantiles of BMI

Continuous outcomes (cont)

321/ 333


APC-model for quantitative outcomes

- ▶ Australian surveys
- ▶ 40,000+ person surveyed at different times
- ▶ Date of birth, data of survey, sex and BMI known.
- ▶ How does BMI evolve **in the population?**
- ▶ Linear model ($E(\text{BMI})$)
- ▶ Quantile regression (median, quantile)
- ▶ — the latter is not a model




Continuous outcomes (cont)

322/ 333



Continuous outcomes (cont)

323/ 333



Continuous outcomes (cont)

324/ 333

Continuous outcomes

Bendix Carstensen

Statistical Analysis in Lexis Diagrams:
Age-Period-Cohort models
— and some cousins
European Doctoral School of Demography,
Centre d'Estudis Demogràfics, Barcelona (virtual), May 25–28 2020

<http://BendixCarstensen.com/APC/EDSD-2020>

cont

