

Analysis of multistate data with realistic rate models and multiple time scales: A dogmatic approach

Bendix Carstensen Steno Diabetes Center
Gentofte, Denmark
<http://BendixCarstensen.com>

IARC, Lyon, France,
11 April 2018

1/ 51

do not count people after they are dead

- ▶ The estimation of the subdistribution hazard boils down to:

$$\tilde{h}(t) = P \{X(t + dt) = j | X(t) \neq j\} / dt$$

that is, the instantaneous rate of failure per time unit from cause j among those who are either alive or have died from causes other than j at time t

- ▶ ... sounds crazy, but...
- ▶ when modeling the **cumulative risk** you must refer back to the size of the **original** population, which include those dead from other causes.
- ▶ The debate is rather if the subdistribution hazard is a useful scale for modeling and reporting from competing risk settings

5/ 51

The dogma [1]

- ▶ do not condition on the future — **indisputable**
- ▶ do not count people after they are dead — **disputable**
- ▶ stick to this world — **expandable**

2/ 51

stick to this world

- ▶ the “net” survival or “cause specific survival”:

$$S_c(t) = \exp \left(\int_0^t \lambda_c(s) ds \right)$$

- ▶ not a proper probability
- ▶ the probability of survival if
 - ▶ all other causes of death than c were absent
 - ▶ c -specific mortality rate were still the same
- ▶ so it is just a transformation of the cause-specific rate with no real world interpretation
- ▶ ... do not label quantities “survival” or “probability” when they are not (of this world)

6/ 51

do not condition on the future

- ▶ commonly seen in connection with “immortal time bias”
- ▶ allocation of follow-up (risk time) to a covariate value only assumed in the future
- ▶ all follow-up among persons **ever** on insulin allocated to the insulin group
 - including the time **prior** to insulin use (when not on insulin)
- ▶ events always with the correct covariate values
- ▶ ⇒ too **much** PY in insulin group; rates too **small**
- ▶ ⇒ too **little** PY in non-insulin group; rates too **large**
- ▶ ⇒ insulin vs. non-insulin rates **underestimated**

3/ 51

(further) dogma for “sticking to this world”

- ▶ rates are continuous in time (and “smooth”)
- ▶ rates may depend on more than one time scale
- ▶ which, is an empirical question

7/ 51

do not count people after they are dead

- ▶ Reference to Fine & Gray’s paper on models for the subdistribution hazard [2]
- ▶ Recall: hazard and cumulative risk for all cause death:

$$F(t) = 1 - \exp(-\Lambda(t)) \Leftrightarrow \lambda(t) = \Lambda'(t) = (\log(1 - F(t)))'$$

- ▶ Subdistribution hazard — with more causes of death (competing risks), for cumulative risk of cause c , $F_c(t)$:

$$\tilde{\lambda}_c(t) = (\log(1 - F_c(t)))'$$

- ▶ Note: F_c depends on all cause-specific hazards

4/ 51

A look at the Cox model

$$\lambda(t, x) = \lambda_0(t) \times \exp(x'\beta)$$

A model for the rate as a function of t and x .

Covariates:

- ▶ x
- ▶ t
- ▶ ... often the effect of t is ignored (forgotten?)
- ▶ *i.e.* left unreported

8/ 51

The Cox-likelihood as profile likelihood


- ▶ One parameter per death time to describe the effect of time (i.e. the chosen timescale).

$$\log(\lambda(t, x_i)) = \log(\lambda_0(t)) + \underbrace{\beta_1 x_{1i} + \dots + \beta_p x_{pi}}_{\eta_i} = \alpha_t + \eta_i$$

- ▶ Profile likelihood:
 - ▶ Derive estimates of α_t as function of data and β s
 - assuming constant rate between death/censoring times
 - ▶ Insert in likelihood, now only a function of data and β s
 - ▶ This turns out to be Cox's partial likelihood
- ▶ Cumulative intensity ($\Lambda_0(t)$) obtained via the Breslow-estimator


9/ 51

Mayo Clinic lung cancer 60 year old woman



13/ 51

Mayo Clinic lung cancer data: 60 year old woman



10/ 51

Example: Mayo Clinic lung cancer I

```
> library( survival )
> library( Epi )
> Lung <- Lexis( exit = list( tfe=time ),
+             exit.status = factor(status,labels=c("Alive","Dead")),
+             data = lung )
```

NOTE: entry.status has been set to "Alive" for all.
NOTE: entry is assumed to be 0 on the tfe timescale.

```
> summary( Lung )
```

```
Transitions:
  To
From Alive Dead Records Events Risk time Persons
  Alive 63 165      228     165  69593      228
```

14/ 51

Splitting the dataset a priori

- ▶ The Poisson approach needs a dataset of empirical rates (d, y) with suitably small values of y .
- ▶ — each individual contributes many empirical rates
- ▶ (one per risk-set contribution in Cox-modelling)
- ▶ From each empirical rate we get:
 - ▶ Poisson-response d
 - ▶ Risk time $y \rightarrow \log(y)$ as offset
 - ▶ time scale covariates: current age, current date, ...
 - ▶ other covariates
- ▶ Contributions not independent, but likelihood is a product
- ▶ Same likelihood as for independent Poisson variates
- ▶ Poisson `glm` with spline/factor effect of time

11/ 51

Example: Mayo Clinic lung cancer II

```
> system.time(
+ mL.cox <- coxph( Surv( tfe, tfe+lex.dur, lex.Xst=="Dead" ) ~
+                 age + factor( sex ),
+                 method="breslow", data=Lung ) )
```

```
user system elapsed
0.010 0.001 0.009
```

```
> Lung.s <- splitLexis( Lung,
+                      breaks=c(0,sort(unique(Lung$time))),
+                      time.scale="tfe" )
> summary( Lung.s )
```

```
Transitions:
  To
From Alive Dead Records Events Risk time Persons
  Alive 19857 165      20022     165  69593      228
```

```
> subset( Lung.s, lex.id==96 )[1:11] ; nlevels( factor( Lung.s$tfe ) )
```

15/ 51

Example: Mayo Clinic lung cancer

- ▶ Survival after lung cancer
- ▶ Covariates:
 - ▶ Age at diagnosis
 - ▶ Sex
 - ▶ Time since diagnosis
- ▶ Cox model
- ▶ Split data:
 - ▶ Poisson model, time as factor
 - ▶ Poisson model, time as spline

12/ 51

Example: Mayo Clinic lung cancer III

```
lex.id tfe lex.dur lex.Cst lex.Xst inst time status age sex ph.ecog
9235 96 0 5 Alive Alive 12 30 2 72 1 2
9236 96 5 6 Alive Alive 12 30 2 72 1 2
9237 96 11 1 Alive Alive 12 30 2 72 1 2
9238 96 12 1 Alive Alive 12 30 2 72 1 2
9239 96 13 2 Alive Alive 12 30 2 72 1 2
9240 96 15 11 Alive Alive 12 30 2 72 1 2
9241 96 26 4 Alive Dead 12 30 2 72 1 2
```

```
[1] 186
```

```
> system.time(
+ mLs.pois.fc <- glm( lex.Xst=="Dead" ~ - 1 + factor( tfe ) +
+                  age + factor( sex ),
+                  offset = log(lex.dur),
+                  family=poisson, data=Lung.s, eps=10^-8, maxit=25 )
+ )
```

```
user system elapsed
13.550 17.334 8.761
```

16/ 51

Example: Mayo Clinic lung cancer IV

```
> length( coef(mLs.pois.fc) )
[1] 188
> t.kn <- c(0,25,100,500,1000)
> dim( Ns(Lung.s$tfe,knots=t.kn) )
[1] 20022 4
> system.time(
+ mLs.pois.sp <- glm( lex.Xst=="Dead" ~ Ns( tfe, knots=t.kn ) +
+                   age + factor( sex ),
+                   offset = log(lex.dur),
+                   family=poisson, data=Lung.s, eps=10^-8, maxit=25 )
+ )
user system elapsed
0.418 0.510 0.317
```

17/ 51

Deriving the survival function

```
> mLs.pois.sp <- glm( lex.Xst=="Dead" ~ Ns( tfe, knots=t.kn ) +
+                   age + factor( sex ),
+                   offset = log(lex.dur),
+                   family=poisson, data=Lung.s, eps=10^-8, maxit=25 )
> CM <- cbind( 1, Ns( seq(10,1000,10)-5, knots=t.kn ), 60, 1 )
> lambda <- ci.exp( mLs.pois.sp, ctr.mat=CM )
> Lambda <- ci.cum( mLs.pois.sp, ctr.mat=CM, int1=10 )[, -4]
> survP <- exp(-rbind(0,Lambda))
```

Code and output for the entire example available in <http://bendixcarstensen.com/AdvCoh/WNtCMA/>

20/ 51

Example: Mayo Clinic lung cancer V

```
> ests <-
+ rbind( ci.exp(mL.cox),
+        ci.exp(mLs.pois.fc,subset=c("age","sex")),
+        ci.exp(mLs.pois.sp,subset=c("age","sex")))
> cmp <- cbind( ests[c(1,3,5),,],
+              ests[c(1,3,5)+1,,] )
> rownames( cmp ) <- c("Cox","Poisson-factor","Poisson-spline")
> colnames( cmp )[c(1,4)] <- c("age","sex")

> round( cmp, 7 )
              age      2.5%      97.5%      sex      2.5%      97.5%
Cox           1.017158 0.9989388 1.035710 0.5989574 0.4313720 0.8316487
Poisson-factor 1.017158 0.9989388 1.035710 0.5989574 0.4313720 0.8316487
Poisson-spline 1.016189 0.9980329 1.034676 0.5998287 0.4319932 0.8328707
```

18/ 51

What the Cox-model really is

Taking the life-table approach *ad absurdum* by:

- ▶ dividing time very finely and
- ▶ modeling one covariate, the time-scale, with one parameter per distinct value.
- ▶ the **model** for the time scale is really with exchangeable time-intervals.
- ▶ ⇒ difficult to access the baseline hazard (which looks terrible)
- ▶ ⇒ uninitiated tempted to show survival curves where irrelevant

21/ 51

Models of this world

- ▶ Replace the α_t s by a parametric function $f(t)$ with a limited number of parameters, for example:
 - ▶ Piecewise constant
 - ▶ Splines (linear, quadratic or cubic)
 - ▶ Fractional polynomials
- ▶ the two latter brings model into "this world":
 - ▶ smoothly varying rates
 - ▶ parametric closed form representation of baseline hazard
 - ▶ finite no. of parameters
- ▶ Makes it really easy to use rates directly in calculations of
 - ▶ expected residual life time
 - ▶ state occupancy probabilities in multistate models
 - ▶ ...


22/ 51

additional dogma


Not sacred, merely consequences of the 3rd commandment:

- ▶ Show risk time **in** states and transitions **between** states graphically
- ▶ Model transition rates by smooth parametric functions
- ▶ There is no such thing as primary or secondary time scale — time scales and other quantitative covariates should be modeled the same way
- ▶ Determine the relevant timescale(s)
- ▶ **Then** derive the relevant measures to report.
- ▶ Time-scale interactions is the proper name for "non-proportional hazards"
- ▶ Multiple time scales should be reported jointly

23/ 51




19/ 51



19/ 51




EBMT transplant data

Iacobelli & Carstensen: Multistate Models with Multiple Timescales, Stat Med 2013, [3]



other covariates: Age and date at Tx, sex, donor type, CML type

24/ 51





Markov property: Empirical question

Model for mortality rates:

- ▶ t time since transplant
- ▶ r time since relapse (if relapsed)
- ▶ t_r time from transplant to relapse
- ▶ Fit the model for all transitions:
 - ▶ split follow-up time
 - ▶ fit Poisson model with covariates
 - ▶ and spline terms for each time scale.
- ▶ Lexis machinery [4, 5] from the Epi package for R
- ▶ ... for representation and manipulation of follow-up data.

26/ 51



Diabetologia
DOI 10.1007/s00125-016-4065-6



ARTICLE





Years of life gained by multifactorial intervention in patients with type 2 diabetes mellitus and microalbuminuria: 21 years follow-up on the Steno-2 randomised trial

Peter Gaede^{1,2} · Jens Oelgaard^{1,2,3} · Bendix Carstensen³ · Peter Rossing^{3,4,5} · Henrik Lund-Andersen^{3,5,6} · Hans-Henrik Parving^{5,7} · Oluf Pedersen⁸

Received: 7 April 2016 / Accepted: 1 July 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract
Aims/hypothesis The aim of this work was to study the potential long-term impact of a 7.8 years intensified, multifactorial pharmacological approaches. After 7.8 years the study continued as an observational follow-up with all patients receiving treatment as for the original intensive therapy group. The pri-

31/ 51





- ### Models used
- ▶ One model for the 4 mortality rates
 - ▶ One model for the 3 CVD rates
 - ▶ ... both models assume:
 - ▶ proportional hazards between CVD states (0, 1, 2, 3) CVD events
 - ▶ proportional hazards between groups (conventional, intervention)
 - ▶ proportional hazards between levels of sex and age (at entry)
 - ▶ Which just means: multiplicative effects of the covariates: **time since baseline**, CVD state, group, sex and age
 - ▶ **Proportional hazards** means: no interaction with the **time scale**
- 34/ 51

Hazard ratios

	Mortality	CVD event
HR, Int. vs. Conv.	0.83 (0.54; 1.30)	0.55 (0.39;0.77)
H ₀ : PH btw. CVD groups	p=0.438	p=0.261
H ₀ : HR = 1	p=0.425	p=0.001
HR vs. 0 CVD events:		
0 (ref.)	1.00	1.00
1	3.08 (1.82; 5.19)	2.43 (1.67;3.52)
2	4.42 (2.36; 8.29)	3.48 (2.15;5.64)
3+	7.76 (4.11;14.65)	

Then use fitted rates to estimate the probabilities of being in each state at all times. (This is immensely complicated).

35/ 51



Expected lifetime and YLL (well, gained)

- Expected lifetime (years) in the Steno 2 cohort during the first 20 years after baseline by treatment group and CVD status.

State	where	Int.	Conv.	Int.–Conv.
Alive	under black line	15.6	14.1	1.5
No CVD	green area	12.7	10.0	2.6
Any CVD	orange area	3.0	4.1	-1.1

- What does "expected" mean?
- Expectation w.r.t. age and sex-distribution in the Steno2 study!
- Computed as areas under survival curves

40 / 51

Multistate models in practice:

- Representation:



- States
- Transitions
- Sojourn times
- Rates

- Analysis of rates:

- Cox-model
- Poisson model

- Reporting

- Rates
- HRs
- Probabilities
- Expected lifetime




41 / 51

From rates to probabilities

- There is a one-to-one correspondence between:
 - all rates between states (by time) + initial state distribution
 - state distribution by time
- Model for rates
 - ⇒ probability of being in a given state at any given time
- Analytically this is a nightmare
- Simulation is the answer

45 / 51



42 / 51

From rates to probabilities: simLexis


- Assume a person is in "DM" initially
 - Simulate a time of death (transition to "D(no CVD)")
 - Simulate a time of CVD (transition to "1st CVD")
 - Choose the smaller as the transition
- If transition is to "1st CVD" simulate death / 2nd CVD, etc.
- Repeat for, say, 10,000 persons
 - ⇒ simulated cohort study
- simLexis does this for you, provided you have
 - initial state and covariates for all persons
 - models to predict (cumulative) rates
- Count how many is in each state at each time:
 - ⇒ state occupancy probabilities
- nState and pState does this for you

46 / 51

Expected lifetime (years) during the first 20 years after baseline by sex, age, treatment group and CVD status.

sex	state	age	Men			Women		
			Int.	Conv.	Int.–Conv.	Int.	Conv.	Int.–Conv.
Alive	45	18.5	17.5	1.0	19.1	18.4	0.7	
	50	17.2	16.1	1.1	18.0	17.2	0.8	
	55	15.6	13.8	1.8	17.4	15.9	1.6	
	60	13.9	11.6	2.2	15.5	13.7	1.8	
	65	11.2	9.5	1.8	13.3	11.4	2.0	
No CVD	45	14.9	12.5	2.4	15.8	14.3	1.5	
	50	14.0	11.1	2.9	15.1	12.9	2.2	
	55	12.2	9.7	2.5	14.3	11.6	2.7	
	60	10.9	8.2	2.7	12.4	9.9	2.6	
	65	9.0	6.7	2.2	10.7	8.3	2.4	

43 / 51



47 / 51

Using the Lexis machinery

- ▶ Allows estimation of fully parametric rate function
- ▶ Simple test for proportional hazards
- ▶ State occupancy probabilities requires simulation: [simLexis](#) — see vignette in [Epi](#) package
- ▶ Access to other measures such as expected residual lifetime.
- ▶ — similar machinery available in Stata:
 - ▶ multistate
 - ▶ Crowther & Lambert [6]
 - ▶ Only one timescale, however...

48/ 51

Additional dogma


- ▶ Show risk time **in** states and transitions **between** states
- ▶ Model transition rates by smooth **parametric** functions
- ▶ There is no such thing as primary or secondary time scale — **time scales** and other quantitative covariates should be modeled the same way
- ▶ Time-scale **interactions** is the proper name for “non-proportional hazards”
- ▶ **Multiple** time scales should be reported jointly

49/ 51

References I

-  P. K. Andersen and N. Keiding. Interpretability and importance of functionals in competing risks and multistate models. *Stat Med*, 31:1074–1088, 2012.
-  J P Fine and R J Gray. A proportional hazards model for the subdistribution of a competing risk. *Journal of the American Statistical Association*, 94(446), 1999.
-  S. Iacobelli and B. Carstensen. Multiple time scales in multi-state models. *Stat Med*, 32(30):5315–5327, Dec 2013.
-  Martyn Plummer and Bendix Carstensen. Lexis: An R class for epidemiological studies with long-term follow-up. *Journal of Statistical Software*, 38(5):1–12, 1 2011.
-  Bendix Carstensen and Martyn Plummer. Using Lexis objects for multi-state models in R. *Journal of Statistical Software*, 38(6):1–18, 1 2011.

References II

-  M. J. Crowther and P. C. Lambert. Parametric multistate survival models: Flexible modelling allowing transition-specific distributions with application to estimating clinically useful measures of effect differences. *Stat Med*, 36(29):4719–4742, Dec 2017.

bendixcarstensen.com/AdvCoh/Lexis-ex